DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY

Base from U.S. Geological Survey
Casa Diablo Mountain, Devils Postpile, Mono Craters, Mount Morrison, 1953.
Bodie, 1958; Cowtrack Mountain, Glass Mountain, 1982
Polyconic projection

SCALE 1:62 500

CONTOUR INTERVALS 40 AND 80 FEET
DOTTED LINES REPRESENT HALF-INTERVAL CONTOURS
NATIONAL GEODETIC VERTICAL DATUM OF 1929

Geology compiled in 1965-1988 by
Quaternary volcanic rocks and
assisted by Robert P. Koeppen
Wilson, Roy E. Chaudet, and Mort
1978, 1981, 1985, 1986, Sierra
generalized from unpublished mapp
1977, Geology of the mountain
generalized from unpublished mapp
1986, and R.A. Bailey, 1974-1975
northern part of the Glass Mou
Krauskopf and Bateman (1977) a
Cowtrack Mountain quadrangle (L
Michael F. Sheridan (1973), Pe
Rinehart and Ross (1957, 1964, 1
(1966), and Krauskopf and Bateman

GEOLOGIC MAP OF LONG VALLEY CALDERA, MONO-INYO CRATERS VOLCANIC CHAIN, AND VICINITY, EASTERN CALIFORNIA

By
Roy A. Bailey

1989

best copy available

MISCELLANEOUS INVESTIGATIONS SERIES
MAP I-1933 SHEET 1 OF 2
Explanatory pamphlet accompanies map

Qal
Qc
Qoa
Qmr
Qbt
Qpa
Qtt
Qgs
Qbs
Qbm
Qbl
Qbo
Qbp
Qbr
Qbu
Qbv
Qbw
Qbx
Qby
Qbz
Qca
Qcb
Qcc
Qcd
Qce
Qcf
Qcg
Qch
Qci
Qcj
Qck
Qcl
Qcm
Qcn
Qco
Qcp
Qcq
Qcr
Qcs
Qct
Qcu
Qcv
Qcw
Qcx
Qcy
Qcz
Qda
Qdb
Qdc
Qdd
Qde
Qdf
Qdg
Qdh
Qdi
Qdj
Qdk
Qdl
Qdm
Qdn
Qdo
Qdp
Qdq
Qdr
Qds
Qdt
Qdu
Qdv
Qdw
Qdx
Qdy
Qdz
Qea
Qeb
Qec
Qed
Qee
Qef
Qeg
Qeh
Qei
Qej
Qek
Qel
Qem
Qen
Qeo
Qep
Qeq
Qer
Qes
Qet
Qeu
Qev
Qew
Qex
Qey
Qez
Qfa
Qfb
Qfc
Qfd
Qfe
Qff
Qfg
Qfh
Qfi
Qfj
Qfk
Qfl
Qfm
Qfn
Qfo
Qfp
Qfq
Qfr
Qfs
Qft
Qfu
Qfv
Qfw
Qfx
Qfy
Qfz
Qga
Qgb
Qgc
Qgd
Qge
Qgf
Qgg
Qgh
Qgi
Qgj
Qgk
Qgl
Qgm
Qgn
Qgo
Qgp
Qgq
Qgr
Qgs
Qgt
Qgu
Qgv
Qgw
Qgx
Qgy
Qgz
Qha
Qhb
Qhc
Qhd
Qhe
Qhf
Qhg
Qhh
Qhi
Qhj
Qhk
Qhl
Qhm
Qhn
Qho
Qhp
Qhq
Qhr
Qhs
Qht
Qhu
Qhv
Qhw
Qhx
Qhy
Qhz
Qia
Qib
Qic
Qid
Qie
Qif
Qig
Qih
Qii
Qij
Qik
Qil
Qim
Qin
Qio
Qip
Qiq
Qir
Qis
Qit
Qiu
Qiv
Qiw
Qix
Qiy
Qiz
Qja
Qjb
Qjc
Qjd
Qje
Qjf
Qjg
Qjh
Qji
Qjj
Qjk
Qjl
Qjm
Qjn
Qjo
Qjp
Qjq
Qjr
Qjs
Qjt
Qju
Qjv
Qjw
Qjx
Qjy
Qjz
Qka
Qkb
Qkc
Qkd
Qke
Qkf
Qkg
Qkh
Qki
Qkj
Qkk
Qkl
Qkm
Qkn
Qko
Qkp
Qkq
Qkr
Qks
Qkt
Qku
Qkv
Qkw
Qkx
Qky
Qkz
Qla
Qlb
Qlc
Qld
Qle
Qlf
Qlg
Qlh
Qli
Qlj
Qlk
Qll
Qlm
Qln
Qlo
Qlp
Qlq
Qlr
Qls
Qlt
Qlu
Qlv
Qlw
Qlx
Qly
Qlz
Qma
Qmb
Qmc
Qmd
Qme
Qmf
Qmg
Qmh
Qmi
Qmj
Qmk
Qml
Qmm
Qmn
Qmo
Qmp
Qmq
Qmr
Qms
Qmt
Qmu
Qmv
Qmw
Qmx
Qmy
Qmz
Qna
Qnb
Qnc
Qnd
Qne
Qnf
Qng
Qnh
Qni
Qnj
Qnk
Qnl
Qnm
Qnn
Qno
Qnp
Qnq
Qnr
Qns
Qnt
Qnu
Qnv
Qnw
Qnx
Qny
Qnz
Qoa
Qob
Qoc
Qod
Qoe
Qof
Qog
Qoh
Qoi
Qoj
Qok
Qol
Qom
Qon
Qoo
Qop
Qoq
Qor
Qos
Qot
Qou
Qov
Qow
Qox
Qoy
Qoz
Qpa
Qpb
Qpc
Qpd
Qpe
Qpf
Qpg
Qph
Qpi
Qpj
Qpk
Qpl
Qpm
Qpn
Qpo
Qpp
Qpq
Qpr
Qps
Qpt
Qpu
Qpv
Qpw
Qpx
Qpy
Qpz
Qqa
Qqb
Qqc
Qqd
Qqe
Qqf
Qqg
Qqh
Qqi
Qqj
Qqk
Qql
Qqm
Qqn
Qqo
Qqp
Qqq
Qqr
Qqs
Qqt
Qqu
Qqv
Qqw
Qqx
Qqy
Qqz
Qra
Qrb
Qrc
Qrd
Qre
Qrf
Qrg
Qrh
Qri
Qrj
Qrk
Qrl
Qrm
Qrn
Qro
Qrp
Qrq
Qrr
Qrs
Qrt
Qru
Qrv
Qrw
Qrx
Qry
Qrz
Qsa
Qsb
Qsc
Qsd
Qse
Qsf
Qsg
Qsh
Qsi
Qsj
Qsk
Qsl
Qsm
Qsn
Qso
Qsp
Qsq
Qsr
Qss
Qst
Qsu
Qsv
Qsw
Qsx
Qsy
Qsz
Qta
Qtb
Qtc
Qtd
Qte
Qtf
Qtg
Qth
Qti
Qtj
Qtk
Qtl
Qtm
Qtn
Qto
Qtp
Qtq
Qtr
Qts
Qtt
Qtu
Qtv
Qtw
Qtx
Qty
Qtz
Qua
Qub
Quc
Qud
Que
Quf
Qug
Quh
Qui
Quj
Quk
Qul
Qum
Qun
Quo
Qup
Quq
Qur
Qus
Qut
Quu
Quv
Quw
Qux
Quy
Quz
Qva
Qvb
Qvc
Qvd
Qve
Qvf
Qvg
Qvh
Qvi
Qvj
Qvk
Qvl
Qvm
Qvn
Qvo
Qvp
Qvq
Qvr
Qvs
Qvt
Qvu
Qvv
Qvw
Qvx
Qvy
Qvz
Qwa
Qwb
Qwc
Qwd
Qwe
Qwf
Qwg
Qwh
Qwi
Qwj
Qwk
Qwl
Qwm
Qwn
Qwo
Qwp
Qwq
Qwr
Qws
Qwt
Qwu
Qwv
Qww
Qwx
Qwy
Qwz
Qxa
Qxb
Qxc
Qxd
Qxe
Qxf
Qxg
Qxh
Qxi
Qxj
Qxk
Qxl
Qxm
Qxn
Qxo
Qxp
Qxq
Qxr
Qxs
Qxt
Qxu
Qxv
Qxw
Qxx
Qxy
Qxz
Qya
Qyb
Qyc
Qyd
Qye
Qyf
Qyg
Qyh
Qyi
Qyj
Qyk
Qyl
Qym
Qyn
Qyo
Qyp
Qyq
Qyr
Qys
Qyt
Qyu
Qyv
Qyw
Qyx
Qyy
Qyz
Qza
Qzb
Qzc
Qzd
Qze
Qzf
Qzg
Qzh
Qzi
Qzj
Qzk
Qzl
Qzm
Qzn
Qzo
Qzp
Qzq
Qzr
Qzs
Qzt
Qzu
Qzv
Qzw
Qzx
Qzy
Qzz
Jtg
Jtm
Jtp
Jtq
Jtr
Jts
Jtt
Jtu
Jtv
Jtw
Jtx
Jty
Jtz
Jua
Jub
Juc
Jud
Jue
Juf
Jug
Juh
Jui
Juj
Juk
Jul
Jum
Jun
Juo
Jup
Juq
Jur
Jus
Jut
Juu
Juv
Juw
Jux
Juy
Juz
Jva
Jvb
Jvc
Jvd
Jve
Jvf
Jvg
Jvh
Jvi
Jvj
Jvk
Jvl
Jvm
Jvn
Jvo
Jvp
Jvq
Jvr
Jvs
Jvt
Jvu
Jvv
Jvw
Jvx
Jvy
Jvz
Jwa
Jwb
Jwc
Jwd
Jwe
Jwf
Jwg
Jwh
Jwi
Jwj
Jwk
Jwl
Jwm
Jwn
Jwo
Jwp
Jwq
Jwr
Jws
Jwt
Jwu
Jwv
Jww
Jwx
Jwy
Jwz
Jxa
Jxb
Jxc
Jxd
Jxe
Jxf
Jxg
Jxh
Jxi
Jxj
Jxk
Jxl
Jxm
Jxn
Jxo
Jxp
Jxq
Jxr
Jxs
Jxt
Jxu
Jxv
Jxw
Jxx
Jxy
Jxz
Jya
Jyb
Jyc
Jyd
Jye
Jyf
Jyg
Jyh
Jyi
Jyj
Jyk
Jyl
Jym
Jyn
Jyo
Jyp
Jyq
Jyr
Jys
Jyt
Jyu
Jyv
Jyw
Jyx
Jyy
Jyz
Jza
Jzb
Jzc
Jzd
Jze
Jzf
Jzg
Jzh
Jzi
Jzj
Jzk
Jzl
Jzm
Jzn
Jzo
Jzp
Jzq
Jzr
Jzs
Jzt
Jzu
Jzv
Jzw
Jzx
Jzy
Jzz
Kga
Kgb
Kgc
Kgd
Kge
Kgf
Kgg
Kgh
Kgi
Kgj
Kgk
Kgl
Kgm
Kgn
Kgo
Kgp
Kgq
Kgr
Kgs
Kgt
Kgu
Kgv
Kgw
Kgx
Kgy
Kgz
Kha
Khb
Khc
Khd
Khe
Khf
Khg
Khh
Khi
Khj
Khk
Khl
Khm
Khn
Kho
Khp
Khq
Khr
Khs
Kht
Khu
Khv
Khw
Khx
Khy
Khz
Kia
Kib
Kic
Kid
Kie
Kif
Kig
Kih
Kii
Kij
Kik
Kil
Kim
Kin
Kio
Kip
Kiq
Kir
Kis
Kit
Kiu
Kiv
Kiw
Kix
Kiy
Kiz
Kja
Kjb
Kjc
Kjd
Kje
Kjf
Kjg
Kjh
Kji
Kjj
Kjk
Kjl
Kjm
Kjn
Kjo
Kjp
Kjq
Kjr
Kjs
Kjt
Kju
Kjv
Kjw
Kjx
Kjy
Kjz
Kka
Kkb
Kkc
Kkd
Kke
Kkf
Kkg
Kkh
Kki
Kkj
Kkk
Kkl
Kkm
Kkn
Kko
Kkp
Kkq
Kkr
Kks
Kkt
Kku
Kkv
Kkw
Kkx
Kky
Kkz
Kla
Klb
Klc
Kld
Kle
Klf
Klg
Klh
Kli
Klj
Klk
Kll
Klm
Kln
Klo
Klp
Klq
Klr
Kls
Klt
Klu
Klv
Klw
Klx
Kly
Klz
Kma
Kmb
Kmc
Kmd
Kme
Kmf
Kmg
Kmh
Kmi
Kmj
Kmk
Kml
Kmm
Kmn
Kmo
Kmp
Kmq
Kmr
Kms
Kmt
Kmu
Kmv
Kmw
Kmx
Kmy
Kmz
Kna
Knb
Knc
Knd
Kne
Knf
Kng
Knh
Kni
Knj
Knk
Knl
Knm
Knn
Kno
Knp
Knq
Knr
Kns
Knt
Knu
Knv
Knw
Knx
Kny
Knz
Koa
Kob
Koc
Kod
Koe
Kof
Kog
Koh
Koi
Koj
Kok
Kol
Kom
Kon
Koo
Kop
Koq
Kor
Kos
Kot
Kou
Kov
Kow
Kox
Koy
Koz
Kpa
Kpb
Kpc
Kpd
Kpe
Kpf
Kpg
Kph
Kpi
Kpj
Kpk
Kpl
Kpm
Kpn
Kpo
Kpp
Kpq
Kpr
Kps
Kpt
Kpu
Kpv
Kpw
Kpx
Kpy
Kpz
Kqa
Kqb
Kqc
Kqd
Kqe
Kqf
Kqg
Kqh
Kqi
Kqj
Kqk
Kql
Kqm
Kqn
Kqo
Kqp
Kqq
Kqr
Kqs
Kqt
Kqu
Kqv
Kqw
Kqx
Kqy
Kqz
Kra
Krb
Krc
Krd
Kre
Krf
Krg
Krh
Kri
Krj
Krk
Krl
Krm
Krn
Kro
Krp
Krq
Krr
Krs
Krt
Kru
Krv
Krw
Krx
Kry
Krz
Ksa
Ksb
Ksc
Ksd
Kse
Ksf
Ksg
Ksh
Ksi
Ksj
Ksk
Ksl
Ksm
Ksn
Kso
Ksp
Ksq
Ksr
Kss
Kst
Ksu
Ksv
Ksw
Ksx
Ksy
Ksz
Kta
Ktb
Ktc
Ktd
Kte
Ktf
Ktg
Kth
Kti
Ktj
Ktk
Ktl
Ktm
Ktn
Kto
Ktp
Ktq
Ktr
Kts
Ktt
Ktu
Ktv
Ktw
Ktx
Kty
Ktz
Kua
Kub
Kuc
Kud
Kue
Kuf
Kug
Kuh
Kui
Kuj
Kuk
Kul
Kum
Kun
Kuo
Kup
Kuq
Kur
Kus
Kut
Kuu
Kuv
Kuw
Kux
Kuy
Kuz
Kva
Kvb
Kvc
Kvd
Kve
Kvf
Kvg
Kvh
Kvi
Kvj
Kvk
Kvl
Kvm
Kvn
Kvo
Kvp
Kvq
Kvr
Kvs
Kvt
Kvu
Kvv
Kvw
Kvx
Kvy
Kvz
Kwa
Kwb
Kwc
Kwd
Kwe
Kwf
Kwg
Kwh
Kwi
Kwj
Kwk
Kwl
Kwm
Kwn
Kwo
Kwp
Kwq
Kwr
Kws
Kwt
Kwu
Kwv
Kww
Kwx
Kwy
Kwz
Kxa
Kxb
Kxc
Kxd
Kxe
Kxf
Kxg
Kxh
Kxi
Kxj
Kxk
Kxl
Kxm
Kxn
Kxo
Kxp
Kxq
Kxr
Kxs
Kxt
Kxu
Kxv
Kxw
Kxx
Kxy
Kxz
Kya
Kyb
Kyc
Kyd
Kye
Kyf
Kyg
Kyh
Kyi
Kyj
Kyk
Kyl
Kym
Kyn
Kyo
Kyp
Kyq
Kyr
Kys
Kyt
Kyu
Kyv
Kyw
Kyx
Kyy
Kyz
Kza
Kzb
Kzc
Kzd
Kze
Kzf
Kzg
Kzh
Kzi
Kzj
Kzk
Kzl
Kzm
Kzn
Kzo
Kzp
Kzq
Kzr
Kzs
Kzt
Kzu
Kzv
Kzw
Kzx
Kzy
Kzz
Lga
Lgb
Lgc
Lgd
Lge
Lgf
Lgg
Lgh
Lgi
Lgj
Lgk
Lgl
Lgm
Lgn
Lgo
Lgp
Lgq
Lgr
Lgs
Lgt
Lgu
Lgv
Lgw
Lgx
Lgy
Lgz
Lha
Lhb
Lhc
Lhd
Lhe
Lhf
Lhg
Lhh
Lhi
Lhj
Lhk
Lhl
Lhm
Lhn
Lho
Lhp
Lhq
Lhr
Lhs
Lht
Lhu
Lhv
Lhw
Lhx
Lhy
Lhz
Lia
Lib
Lic
Lid
Lie
Lif
Lig
Lih
Lii
Lij
Lik
Lil
Lim
Lin
Lio
Lip
Liq
Lir
Lis
Lit
Liu
Liv
Liw
Lix
Liy
Liz
Lja
Ljb
Ljc
Ljd
Lje
Ljf
Ljg
Ljh
Lji
Ljj
Ljk
Ljl
Ljm
Ljn
Ljo
Ljp
Ljq
Ljr
Ljs
Ljt
Lju
Ljv
Ljw
Ljx
Ljy
Ljz
Lka
Lkb
Lkc
Lkd
Lke
Lkf
Lkg
Lkh
Lki
Lkj
Lkk
Lkl
Lkm
Lkn
Lko
Lkp
Lkq
Lkr
Lks
Lkt
Lku
Lkv
Lkw
Lkx
Lky
Lkz
Lla
Llb
Llc
Lld
Lle
Llf
Llg
Llh
Lli
Llj
Llk
Lll
Llm
Lln
Llo
Llp
Llq
Llr
Lls
Llt
Llu
Llv
Llw
Llx
Lly
Llz
Lma
Lmb
Lmc
Lmd
Lme
Lmf
Lmg
Lmh
Lmi
Lmj
Lmk
Lml
Lmm
Lmn
Lmo
Lmp
Lmq
Lmr
Lms
Lmt
Lmu
Lmv
Lmw
Lmx
Lmy
Lmz
Lna
Lnb
Lnc
Lnd
Lne
Lnf
Lng
Lnh
Lni
Lnj
Lnk
Lnl
Lnm
Lnn
Lno
Lnp
Lnq
Lnr
Lns
Lnt
Lnu
Lnv
Lnw
Lnx
Lny
Lnz
Loa
Lob
Loc
Lod
Loe
Lof
Log
Loh
Loi
Loj
Lok
Lol
Lom
Lon
Loo
Lop
Loq
Lor
Los
Lot
Lou
Lov
Low
Lox
Loy
Loz
Lpa
Lpb
Lpc
Lpd
Lpe
Lpf
Lpg
Lph
Lpi
Lpj
Lpk
Lpl
Lpm
Lpn
Lpo
Lpp
Lpq
Lpr
Lps
Lpt
Lpu
Lpv
Lpw
Lpx
Lpy
Lpz
Lqa
Lqb
Lqc
Lqd
Lqe
Lqf
Lqg
Lqh
Lqi
Lqj
Lqk
Lql
Lqm
Lqn
Lqo
Lqp
Lqq
Lqr
Lqs
Lqt
Lqu
Lqv
Lqw
Lqx
Lqy
Lqz
Lra
Lrb
Lrc
Lrd
Lre
Lrf
Lrg
Lrh
Lri
Lrj
Lrk
Lrl
Lrm
Lrn
Lro
Lrp
Lrq
Lrr
Lrs
Lrt
Lru
Lrv
Lrw
Lrx
Lry
Lrz
Lsa
Lsb
Lsc
Lsd
Lse
Lsf
Lsg
Lsh
Lsi
Lsj
Lsk
Lsl
Lsm
Lsn
Lso
Lsp
Lsq
Lsr
Lss
Lst
Lsu
Lsv
Lsw
Lsx
Lsy
Lsz
Lta
Ltb
Ltc
Ltd
Lte
Ltf
Ltg
Lth
Lti
Ltj
Ltk
Ltl
Ltm
Ltn
Lto
Ltp
Ltq
Ltr
Lts
Ltt
Ltu
Ltv
Ltw
Ltx
Lty
Ltz
Lua
Lub
Luc
Lud
Lue
Luf
Lug
Luh
Lui
Luj
Luk
Lul
Lum
Lun
Luo
Lup
Luq
Lur
Lus
Lut
Luu
Luv
Luw
Lux
Luy
Luz
Lva
Lvb
Lvc
Lvd
Lve
Lvf
Lvg
Lvh
Lvi
Lvj
Lvk
Lvl
Lvm
Lvn
Lvo
Lvp
Lvq
Lvr
Lvs
Lvt
Lvu
Lvv
Lvw
Lvx
Lvy
Lvz
Lwa
Lwb
Lwc
Lwd
Lwe
Lwf
Lwg
Lwh
Lwi
Lwj
Lwk
Lwl
Lwm
Lwn
Lwo
Lwp
Lwq
Lwr
Lws
Lwt
Lwu
Lwv
Lww
Lwx
Lwy
Lwz
Lxa
Lxb
Lxc
Lxd
Lxe
Lxf
Lxg
Lxh
Lxi
Lxj
Lxk
Lxl
Lxm
Lxn
Lxo
Lxp
Lxq
Lxr
Lxs
Lxt
Lxu
Lxv
Lxw
Lxx
Lxy
Lxz
Lya
Lyb
Lyc
Lyd
Lye
Lyf
Lyg
Lyh
Lyi
Lyj
Lyk
Lyl
Lym
Lyn
Lyo
Lyp
Lyq
Lyr
Lys
Lyt
Lyu
Lyv
Lyw
Lyx
Lyy
Lyz
Lza
Lzb
Lzc
Lzd
Lze
Lzf
Lzg
Lzh
Lzi
Lzj
Lzk
Lzl
Lzm
Lzn
Lzo
Lzp
Lzq
Lzr
Lzs
Lzt
Lzu
Lzv
Lzw
Lzx
Lzy
Lzz
Mga
Mgb
Mgc
Mgd
Mge
Mgf
Mgg
Mgh
Mgi
Mgj
Mgk
Mgl
Mgm
Mgn
Mgo
Mgp
Mgq
Mgr
Mgs
Mgt
Mgu
Mgv
Mgw
Mgx
Mgy
Mgz
Mha
Mhb
Mhc
Mhd
Mhe
Mhf
Mhg
Mhh
Mhi
Mhj
Mhk
Mhl
Mhm
Mhn
Mho
Mhp
Mhq
Mhr
Mhs
Mht
Mhu
Mhv
Mhw
Mhx
Mhy
Mhz
Mia
Mib
Mic
Mid
Mie
Mif
Mig
Mih
Mii
Mij
Mik
Mil
Mim
Min
Mio
Mip
Miq
Mir
Mis
Mit
Miu
Miv
Miw
Mix
Miy
Miz
Mja
Mjb
Mjc
Mjd
Mje
Mjf
Mjg
Mjh
Mji
Mjj
Mjk
Mjl
Mjm
Mjn
Mjo
Mjp
Mjq
Mjr
Mjs
Mjt
Mju
Mjv
Mjw
Mjx
Mjy
Mjz
Mka
Mkb
Mkc
Mkd
Mke
Mkf
Mkg
Mkh
Mki
Mkj
Mkk
Mkl
Mkm
Mkn
Mko
Mkp
Mkq
Mkr
Mks
Mkt
Mku
Mkv
Mkw
Mkx
Mky
Mkz
Mla
Mlb
Mlc
Mld
Mle
Mlf
Mlg
Mlh
Mli
Mlj
Mlk
Mll
Mlm
Mln
Mlo
Mlp
Mlq
Mlr
Mls
Mlt
Mlu
Mlv
Mlw
Mlx
Mly
Mlz
Mma
Mmb
Mmc
Mmd
Mme
Mmf
Mmg
Mmh
Mmi
Mmj
Mmk
Mml
Mmm
Mmn
Mmo
Mmp
Mmq
Mmr
Mms
Mmt
Mmu
Mmv
Mmw
Mmx
Mmy
Mmz
Mna
Mnb
Mnc
Mnd
Mne
Mnf
Mng
Mnh
Mni
Mnj
Mnk
Mnl
Mnm
Mnn
Mno
Mnp
Mnq
Mnr
Mns
Mnt
Mnu
Mnv
Mnw
Mnx
Mny
Mnz
Moa
Mob
Moc
Mod
Moe
Mof
Mog
Moh
Moi
Moj
Mok
Mol
Mom
Mon
Moo
Mop
Moq
Mor
Mos
Mot
Mou
Mov
Mow
Mox
Moy
Moz
Mpa
Mpb
Mpc
Mpd
Mpe
Mpf
Mpg
Mph
Mpi
Mpj
Mpk
Mpl
Mpm
Mpn
Mpo
Mpp
Mpq
Mpr
Mps
Mpt
Mpu
Mpv
Mpw
Mpx
Mpy
Mpz
Mqa
Mqb
Mqc
Mqd
Mqe
Mqf
Mqg
Mqh
Mqi
Mqj
Mqk
Mql
Mqm
Mqn
Mqo
Mqp
Mqq
Mqr
Mqs
Mqt
Mqu
Mqv
Mqw
Mqx
Mqy
Mqz
Mra
Mrb
Mrc
Mrd
Mre
Mrf
Mrg
Mrh
Mri
Mrj
Mrk
Mrl
Mrm
Mrn
Mro
Mrp
Mrq
Mrr
Mrs
Mrt
Mru
Mrv
Mrw
Mrx
Mry
Mrz
Msa
Msb
Msc
Msd
Mse
Msf
Msg
Msh
Msi
Msj
Msk
Msl
Msm
Msn
Mso
Msp
Msq
Msr
Mss
Mst
Msu
Msv
Msw
Msx
Msy
Msz
Mta
Mtb
Mtc
Mtd
Mte
Mtf
Mtg
Mth
Mti
Mtj
Mtk
Mtl
Mtm
Mtn
Mto
Mtp
Mtq
Mtr
Mts
Mtt
Mtu
Mtv
Mtw
Mtx
Mty
Mtz
Mua
Mub
Muc
Mud
Mue
Muf
Mug
Muh
Mui
Muj
Muk
Mul
Mum
Mun
Muo
Mup
Muq
Mur
Mus
Mut
Muu
Muv
Muw
Mux
Muy
Muz
Mva
Mvb
Mvc
Mvd
Mve
Mvf
Mvg
Mvh
Mvi
Mvj
Mvk
Mvl
Mvm
Mvn
Mvo
Mvp
Mvq
Mvr
Mvs
Mvt
Mvu
Mvv
Mvw
Mvx
Mvy
Mvz
Mwa
Mwb
Mwc
Mwd
Mwe
Mwf
Mwg
Mwh
Mwi
Mwj
Mwk
Mwl
Mwm
Mwn
Mwo
Mwp
Mwq
Mwr
Mws
Mwt
Mwu
Mwv
Mww
Mwx
Mwy
Mwz
Mxa
Mxb
Mxc
Mxd
Mxe
Mxf
Mxg
Mxh
Mxi
Mxj
Mxk
Mxl
Mxm
Mxn
Mxo
Mxp
Mxq
Mxr
Mxs
Mxt
Mxu
Mxv
Mxw
Mxx
Mxy
Mxz
Mya
Myb
Myc
Myd
Mye
Myf
Myg
Myh
Myi
Myj
Myk
Myl
Mym
Myn
Myo
Myp
Myq
Myr
Mys
Myt
Myu
Myv
Myw
Myx
Myy
Myz
Mza
Mzb
Mzc
Mzd
Mze
Mzf
Mzg
Mzh
Mzi
Mzj
Mzk
Mzl
Mzm
Mzn
Mzo
Mzp
Mzq
Mzr
Mzs
Mzt
Mzu
Mzv
Mzw
Mzx
Mzy
Mzz
Nga
Ngb
Ngc
Ngd
Nge
Ngf
Ngg
Ngh
Ngi
Ngj
Ngk
Ngl
Ngm
Ngn
Ngo
Ngp
Ngq
Ngr
Ngs
Ngt
Ngu
Ngv
Ngw
Ngx
Ngy
Ngz
Nha
Nhb
Nhc
Nhd
Nhe
Nhf
Nhg
Nhh
Nhi
Nhj
Nhk
Nhl
Nhm
Nhn
Nho
Nhp
Nhq
Nhr
Nhs
Nht
Nhu
Nhv
Nhw
Nhx
Nhy
Nhz
Nia
Nib
Nic
Nid
Nie
Nif
Nig
Nih
Nii
Nij
Nik
Nil
Nim
Nin
Nio
Nip
Niq
Nir
Nis
Nit
Niu
Niv
Niw
Nix
Niy
Niz
Nja
Njb
Njc
Njd
Nje
Njf
Njg
Njh
Nji
Njj
Njk
Njl
Njm
Njn
Njo
Njp
Njq
Njr
Njs
Njt
Nju
Njv
Njw
Njx
Njy
Njz
Nka
Nkb
Nkc
Nkd
Nke
Nkf
Nkg
Nkh
Nki
Nkj
Nkk
Nkl
Nkm
Nkn
Nko
Nkp
Nkq
Nkr
Nks
Nkt
Nku
Nkv
Nkw
Nkx
Nky
Nkz
Nla
Nlb
Nlc
Nld
Nle
Nlf
Nlg
Nlh
Nli
Nlj
Nlk
Nll
Nlm
Nln
Nlo
Nlp
Nlq
Nlr
Nls
Nlt
Nlu
Nlv
Nlw
Nlx
Nly
Nlz
Nma
Nmb
Nmc
Nmd
Nme
Nmf
Nmg
Nmh
Nmi
Nmj
Nmk
Nml
Nmm
Nmn
Nmo
Nmp
Nmq
Nmr
Nms
Nmt
Nmu
Nmv
Nmw
Nmx
Nmy
Nmz
Nna
Nnb
Nnc
Nnd
Nne
Nnf
Nng
Nnh
Nni
Nnj
Nnk
Nnl
Nnm
Nnn
Nno
Nnp
Nnq
Nnr
Nns
Nnt
Nnu
Nnv
Nnw
Nnx
Nny
Nnz
Noa
Nob
Noc
Nod
Noe
Nof
Nog
Noh
Noi
Noj
Nok
Nol
Nom
Non
Noo
Nop
Noq
Nor
Nos
Not
Nou
Nov
Now
Nox
Noy
Noz
Npa
Npb
Npc
Npd
Npe
Npf
Npg
Nph
Npi
Npj
Npk
Npl
Npm
Npn
Npo
Npp
Npq
Npr
Nps
Npt
Npu
Npv
Npw
Npx
Npy
Npz
Nqa
Nqb
Nqc
Nqd
Nqe
Nqf
Nqg
Nqh
Nqi
Nqj
Nqk
Nql
Nqm
Nqn
Nqo
Nqp
Nqq
Nqr
Nqs
Nqt
Nqu
Nqv
Nqw
Nqx
Nqy
Nqz
Nra
Nrb
Nrc
Nrd
Nre
Nrf
Nrg
Nrh
Nri
Nrj
Nrk
Nrl
Nrm
Nrn
Nro
Nrp
Nrq
Nrr
Nrs
Nrt
Nru
Nrv
Nrw
Nrx
Nry
Nrz
Nsa
Nsb
Nsc
Nsd
Nse
Nsf
Nsg
Nsh
Nsi
Nsj
Nsk
Nsl
Nsm
Nsn
Nso
Nsp
Nsq
Nsr
Nss
Nst
Nsu
Nsv
Nsw
Nsx
Nsy
Nsz
Nta
Ntb
Ntc
Ntd
Nte
Ntf
Ntg
Nth
Nti
Ntj
Ntk
Ntl
Ntm
Ntn
Nto
Ntp
Ntq
Ntr
Nts
Ntt
Ntu
Ntv
Ntw
Ntx
Nty
Ntz
Nua
Nub
Nuc
Nud
Nue
Nuf
Nug
Nuh
Nui
Nuj
Nuk
Nul
Num
Nun
Nuo
Nup
Nuq
Nur
Nus
Nut
Nuu
Nuv
Nuw
Nux
Nuy
Nuz
Nva
Nvb
Nvc
Nvd
Nve
Nvf
Nvg
Nvh
Nvi
Nvj
Nvk
Nvl
Nvm
Nvn
Nvo
Nvp
Nvq
Nvr
Nvs
Nvt
Nvu
Nvv
Nvw
Nvx
Nvy
Nvz
Nwa
Nwb
Nwc
Nwd
Nwe
Nwf
Nwg
Nwh
Nwi
Nwj
Nwk
Nwl
Nwm
Nwn
Nwo
Nwp
Nwq
Nwr
Nws
Nwt
Nwu
Nwv
Nww
Nwx
Nwy
Nwz
Nxa
Nxb
Nxc
Nxd
Nxe
Nxf
Nxg
Nxh
Nxi
Nxj
Nxk
Nxl
Nxm
Nxn
Nxo
Nxp
Nxq
Nxr
Nxs
Nxt
Nxu
Nxv
Nxw
Nxx
Nxy
Nxz
Nya
Nyb
Nyc
Nyd
Nye
Nyf
Nyg
Nyh
Nyi
Nyj
Nyk
Nyl
Nym
Nyn
Nyo
Nyp
Nyq
Nyr
Nys
Nyt
Nyu
Nyv
Nyw
Nyx
Nyy
Nyz
Nza
Nzb
Nzc
Nzd
Nze
Nzf
Nzg
Nzh
Nzi
Nzj
Nzk
Nzl
Nzm
Nzn
Nzo
Nzp
Nzq
Nzr
Nzs
Nzt
Nzu
Nzv
Nzw
Nzx
Nzy
Nzz
Oga
Ogb
Ogc
Ogd
Oge
Ogf
Ogg
Ogh
Ogi
Ogj
Ogk
Ogl
Ogm
Ogn
Ogo
Ogp
Ogq
Ogr
Ogs
Ogt
Ogu
Ogv
Ogw
Ogx
Ogy
Ogz
Oha
Ohb
Ohc
Ohd
Ohe
Ohf
Ohg
Ohh
Ohi
Ohj
Ohk
Ohl
Ohm
Ohn
Oho
Ohp
Ohq
Ohr
Ohs
Oht
Ohu
Ohv
Ohw
Ohx
Ohy
Ohz
Oia
Oib
Oic
Oid
Oie
Oif
Oig
Oih
Oii
Oij
Oik
Oil
Oim
Oin
Oio
Oip
Oiq
Oir
Ois
Oit
Oiu
Oiv
Oiw
Oix
Oiy
Oiz
Oja
Ojb
Ojc
Ojd
Oje
Ojf
Ojg
Ojh
Oji
Ojj
Ojk
Ojl
Ojm
Ojn
Ojo
Ojp
Ojq
Ojr
Ojs
Ojt
Oju
Ojv
Ojw
Ojx
Ojy
Ojz
Oka
Okb
Okc
Okd
Oke
Okf
Okg
Okh
Oki
Okj
Okk
Okl
Okm
Okn
Oko
Okp
Okq
Okr
Oks
Okt
Oku
Okv
Okw
Okx
Oky
Okz
Ola
Olb
Olc
Old
Ole
Olf
Olg
Olh
Oli
Olj
Olk
Oll
Olm
Oln
Olo
Olp
Olq
Olr
Ols
Olt
Olu
Olv
Olw
Olx
Oly
Olz
Oma
Omb
Omc
Omd
Ome
Omf
Omg
Omh
Omi
Omj
Omk
Oml
Omm
Omn
Omo
Omp
Omq
Omr
Oms
Omt
Omu
Omv
Omw
Omx
Omy
Omz
Ona
Onb
Onc
Ond
One
Onf
Ong
Onh
Oni
Onj
Onk
Onl
Onm
Onn
Ono
Onp
Onq
Onr
Ons
Ont
Onu
Onv
Onw
Onx
Ony
Onz
Ooa
Oob
Ooc
Ood
Ooe
Oof
Oog
Ooh
Ooi
Ooj
Ook
Ool
Oom
Oon
Ooo
Oop
Ooq
Oor
Oos
Oot
Oou
Oov
Oow
Oox
Ooy
Ooz
Opa
Opb
Opc
Opd
Ope
Opf
Opg
Oph
Opi
Opj
Opk
Opl
Opm
Opn
Opo
Opp
Opq
Opr
Ops
Opt
Opu
Opv
Opw
Opx
Opy
Opz
Oqa
Oqb
Oqc
Oqd
Oqe
Oqf
Oqg
Oqh
Oqi
Oqj
Oqk
Oql
Oqm
Oqn
Oqo
Oqp
Oqq
Oqr
Oqs
Oqt
Oqu
Oqv
Oqw
Oqx
Oqy
Oqz
Ora
Orb
Orc
Ord
Ore
Orf
Org
Orh
Ori
Orj
Ork
Orl
Orm
Orn
Oro
Orp
Orq
Orr
Ors
Ort
Oru
Orv
Orw
Orx
Ory
Orz
Osa
Osb
Osc
Osd
Ose
Osf
Osg
Osh
Osi
Osj
Osk
Osl
Osm
Osn
Oso
Osp
Osq
Osr
Oss
Ost
Osu
Osv
Osw
Osx
Osy
Osz
Ota
Otb
Otc
Otd
Ote
Otf
Otg
Oth
Oti
Otj
Otk
Otl
Otm
Otn
Oto
Otp
Otq
Otr
Ots
Ott
Otu
Otv
Otw
Otx
Oty
Otz
Oua
Oub
Ouc
Oud
Oue
Ouf
Oug
Ouh
Oui
Ouj
Ouk
Oul
Oum
Oun
Ouo
Oup
Ouq
Our
Ous
Out
Ouu
Ouv
Ouw
Oux
Ouy
Ouz
Ova
Ovb
Ovc
Ovd
Ove
Ovf
Ovg
Ovh
Ovi
Ovj
Ovk
Ovl
Ovm
Ovn
Ovo
Ovp
Ovq
Ovr
Ovs
Ovt
Ovu
Ovv
Ovw
Ovx
Ovy
Ovz
Owa
Owb
Owc
Owd
Owe
Owf
Owg
Owh
Owi
Owj
Owk
Owl
Owm
Own
Owo
Owp
Owq
Owr
Ows
Owt
Owu
Owv
Oww
Owx
Owy
Owz
Oxa
Oxb
Oxc
Oxd
Oxe
Oxf
Oxg
Oxh
Oxi
Oxj
Oxk
Oxl
Oxm
Oxn
Oxo
Oxp
Oxq
Oxr
Oxs
Oxt
Oxu
Oxv
Oxw
Oxx
Oxy
Oxz
Oya
Oyb
Oyc
Oyd
Oye
Oyf
Oyg
Oyh
Oyi
Oyj
Oyk
Oyl
Oym
Oyn
Oyo
Oyp
Oyq
Oyr
Oys
Oyt
Oyu
Oyv
Oyw
Oyx
Oyy
Oyz
Oza
Ozb
Ozc
Ozd
Oze
Ozf
Ozg
Ozh
Ozi
Ozj
Ozk
Ozl
Ozm
Ozn
Ozo
Ozp
Ozq
Ozr
Ozs
Ozt
Ozu
Ozv
Ozw
Ozx
Ozy
Ozz
Pga
Pgb
Pgc
Pgd
Pge
Pgf
Pgg
P

DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY



EXPLANATION

Depth in feet
See sheet 1 for Description and Correlation of map units.



INDEX MAP SHOWING 15-MINUTE QUADRANGLES COVERING THE STUDY AREA

Base from U.S. Geological Survey 1:62,500
Mono Craters, 1953; Bodie, Trench Canyon, 1958; Cowtrack Mountain, 1962
Polyconic projection
Mono Lake bathymetry and shoreline from Los Angeles Department of Water and Power, Report No. 15, 1987

SCALE 1:62 500

CONTOUR INTERVALS 40 AND 80 FEET
DOTTED LINES REPRESENT HALF INTERVAL CONTOURS
NATIONAL GEODETIC VERTICAL DATUM OF 1929
(SEE EXPLANATION FOR BATHYMETRY INTERVAL)

Geology compiled in 1987. Geologic mapping of Quaternary volcanic rocks and surficial deposits by Roy A. Bailey, 1975, 1978, 1987. Pre-Quaternary geology generalized from Kistler (1966b) and Chesterman and Gray (1975)

MONO LAKE GEOLOGIC MAP













EXPLANATION

Depth in feet
See sheet 1 for Description and Correlation of map units.



INDEX MAP SHOWING 15-MINUTE QUADRANGLES COVERING THE STUDY AREA



SCALE 1:250 000

LANDSAT SATELLITE IMAGE OF LONG VALLEY CALDERA AND VICINITY—Adapted from the Mariposa, Calif.-Nev. 1° by 2° quadrangle, experimental edition, 1983, Universal Transverse Mercator projection. Imagery recorded by Multispectral Scanner (MSS) on National Aeronautics and Space Administration (NASA) Landsat 4 satellite, on October 26 and November 2, 1983. Rectangles outline area of the Long Valley geologic map and the southern part of the offset Mono Lake geologic map. Colors generally relate to typical features as follows: black, deep clear water; reddish orange, marsh and wet grassland; reddish brown, conifer forests; medium blue, Holocene rhyolite lava; light blue, barren ash-covered land; green, sage; yellow, barren reddish outcrop; white, snow; bright red, irrigated fields.












EXPLANATION
Geologic units—Dark color, area of outcrop; light color, inferred subsurface distribution
Cretaceous granitic rocks
Jurassic granitic rocks
Triassic plutonic rocks
Mesozoic metavolcanic rocks
Paleozoic metasedimentary rocks
Contact
Inferred subsurface contact
Drill hole—Showing subsurface pre-Tertiary bedrock
Mono Craters tunnel—Showing pre-Tertiary bedrock (Gronseth, 1940). Tertiary rock indicated uncolored
Basement lithology uncertain (?)
MONO LAKE
Outline of Long Valley caldera
0 5 10 15 20 KILOMETERS

MAP A—GENERAL DISTRIBUTION OF PRE-TERTIARY ROCKS



MAP B—MAJOR FAULTS AND STRUCTURAL TRENDS





Base from U.S. Geological Survey 1:62 500
Mono Craters, 1953; Blake, Tranch Canyon, 1958; Cowtrack Mountain, 1962
Polyconic projection
Mono Lake bathymetry and shoreline from Los Angeles Department of Water and Power, Report No. 15, 1987

SCALE 1:62 500

CONTOUR INTERVALS 40 AND 80 FEET

Geology compiled in 1987. Geologic mapping of Quaternary volcanic rocks and surficial deposits by Roy A. Bailey, 1975, 1978, 1987. Pre-Quaternary geology generalized from Kistler (1966b) and Chesterman and Gray (1975)



INDEX MAP SHOWING 15-MINUTE QUADRANGLES COVERING THE STUDY AREA

MONO LAKE GEOLOGIC MAP











CROSS SECTIONS



MAP A—GENERAL E

MAP C—PRE

GEOLOGIC MAP OF LONG VALLEY CALDERA, MONO-INYO CRATERS VOLCANIC CHAIN, AND VICINITY, EASTERN CALIFOR

By
Roy A. Bailey

1989

best copy available



Geologic mapping of Quaternary volcanic rocks and Bailey, 1975, 1976, 1981. Pre-Quaternary geology (Kistler and Chesterman and Gray (1975)



INDEX MAP SHOWING 15-MINUTE QUADRANGLES COVERING THE STUDY AREA



LANDSAT SATELLITE IMAGE OF LONG VALLEY CALDERA AND VICINITY—Adapted from the Mariposa, Calif.-Nev. 1° by 2° quadrangle, experimental edition, 1983, Universal Transverse Mercator projection. Imagery recorded by Multispectral Scanner (MSS) on National Aeronautics and Space Administration (NASA) Landsat 4 satellite, on October 26 and November 2, 1983. Rectangles outline area of the Long Valley geologic map and the southern part of the offset Mono Lake geologic map. Colors generally relate to typical features as follows: black, deep clear water; reddish orange, marsh and wet grassland; reddish brown, conifer forests; medium blue, Holocene rhyolite lava; light blue, barren ash-covered land; green, sage; yellow, barren reddish outcrop; white, snow; bright red, irrigated fields.













MAP A—GENERAL DISTRIBUTION OF PRE-TERTIARY ROCKS


EXPLANATION
Topographic margin of Long Valley caldera floor
Topographic margin of resurgent dome
Normal fault—Ball and bar on downthrown side
Major range-bounding fault—Ball and bar on downthrown side
MONO LAKE
SIERRA NEVADA
HARTLEY SPRINGS FAULT
BLACK MOUNTAIN FAULT
BENTON RANGE FAULT ZONE
LONG VALLEY FAULT
WATTERSON CANYON FAULT
CLOVER PATCH FAULT
WILSON CREEK FAULT
Outline of Long Valley caldera
0 5 10 15 20 KILOMETERS

MAP B—MAJOR FAULTS AND STRUCTURAL TRENDS



MAP C—PRECALDERA VOLCANIC VENTS



MAP D—POSTCALDERA VOLCANIC VENTS

OLOGIC MAP OF LONG VALLEY CALDERA, MONO-INYO CRATERS VOLCANIC CHAIN, AND VICINITY, EASTERN CALIFORNIA

By
Roy A. Bailey

1989

best copy available


MAP I-1933
DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY
GEOLOGIC MAP OF THE LONG VALLEY CALDERA,
MONO-INYO CRATERS VOLCANIC CHAIN, AND
VICINITY, EASTERN CALIFORNIA
By Roy A. Bailey
MISCELLANEOUS INVESTIGATIONS SERIES
Published by the U.S. Geological Survey, 1989
G
SHEET--GEOLOGIC MAP OF THE LONG VALLEY CALDERA, MONO-INYO CRATERS VOLCANIC CHAIN, EASTERN CALIFORNIA 1:62,500 MAP I-1933

best copy available

DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY

TO ACCOMPANY MAP I–1933

GEOLOGIC MAP OF LONG VALLEY CALDERA, MONO-INYO CRATERS VOLCANIC CHAIN, AND VICINITY, EASTERN CALIFORNIA

By Roy A. Bailey

GEOLOGIC SETTING

Long Valley caldera and the Mono-Inyo Craters volcanic chain compose a late Tertiary to Quaternary volcanic complex on the west edge of the Basin and Range Province at the base of the Sierra Nevada frontal fault escarpment. The caldera, an east-west-elongate, oval depression 17 by 32 km, is located just northwest of the northern end of the Owens Valley rift and forms a reentrant or offset in the Sierran escarpment, commonly referred to as the "Mammoth embayment." The Mono-Inyo Craters volcanic chain forms a north-trending zone of volcanic vents extending 45 km from the west moat of the caldera to Mono Lake. The prevolcanic basement in the area is mainly Mesozoic granitic rock of the Sierra Nevada batholith and Paleozoic metasedimentary and Mesozoic metavolcanic rocks of the Mount Morrison, Gull Lake, and Ritter Range roof pendants (map A).

PHYSIOGRAPHY

The oval floor of Long Valley caldera ranges in elevation from about 2,000 m in its eastern half, which is dominated by Lake Crowley and the surrounding grass- and sage-covered surface of Long Valley proper, to about 2,600 m in its western half, which is hillier and mostly forested. The topographic walls of the caldera rise steeply to elevations of 3,000–3,600 m, except on the east and southeast, where the rim is a 100-200-m subdued escarpment that forms the northwest edge of the Volcanic Tableland. The highest hills within the central and western parts of the caldera rise to a maximum elevation of 2,800 m. Between the central hills and the caldera walls is an annular depression, referred to as the caldera moat, which is drained on the north and east by Deadman Creek and the Owens River and on the south by Mammoth Creek and its lower reach known as Hot Creek.

Only the northern part of the Mono-Inyo Craters volcanic chain, the Mono Craters proper, forms a prominent physiographic feature—a 17-km-long arcuate ridge, extending southward from Mono Lake to near June Lake. Composed of 30 or more overlapping volcanic domes and craters, the ridge rises about 610 m above the surrounding terrain to a maximum elevation of 2,800 m. Viewed from the west this ridge forms a striking skyline of barren crags mantled with a pale-gray blanket of tephra. The Inyo Craters chain forms a less prominent, discontinuous line of volcanic domes and craters extending 10 km south of the Mono Craters into the west moat of Long Valley caldera.

VOLCANISM

Long Valley caldera

Volcanism in the Long Valley area (Bailey and others, 1976; Bailey, 1982b) began about 3.6 Ma with widespread eruption of trachybasaltic-trachyandesitic lavas on a moderately well dissected upland surface (Huber, 1981). Erosional remnants of these mafic lavas are scattered over a 4,000-km^2 area extending from the Adobe Hills (5–10 km notheast of the map area), around the periphery of Long Valley caldera, and southwestward into the High Sierra. Although these lavas never formed a continuous cover over this region, their wide distribution suggests an extensive mantle source for these initial mafic eruptions. Between 3.0 and 2.5 Ma quartz-latite domes and flows erupted near the north and northwest rims of the present caldera, at and near Bald Mountain and on San Joaquin Ridge (Chaudet, 1986). These intermediate-composition lavas probably represent the onset of deep-crustal magmatic accumulation and differentiation that eventually culminated in formation of the large shallow Long Valley magma chamber, from which subsequent more silicic eruptions originated. The first of these more silicic eruptions occurred in the vicinity of Glass Mountain, on the northeast rim of the present caldera, where a 1,000-m-thick complex of high-silica rhyolite domes, flows, and tuffs accumulated between 2.1 and 0.8 Ma (Metz and Mahood, 1985). The high, central part of this complex, consisting of overlapping and juxtaposed domes, flows, and intrusions, was surrounded by an extensive apron of tuffs, consisting of tephra, small-volume ash-flows, and block-and-ash flows, now preserved only on the north and southeast flanks of the complex. The older of these Glass Mountain rhyolites are somewhat heterogeneous chemically, but those 1.1 Ma and younger in age are quite homogeneous and more voluminous, suggesting that early isolated source bodies did not coalesce to form a single large chamber until the latter part of the Glass Mountain episode (Metz and Mahood, 1985).

At 0.73 Ma, catastrophic rupturing of the roof of the magma chamber resulted in the expulsion of 600 km^3 of rhyolite magma, mainly as hot incandescent ash flows. This partial emptying of the chamber caused collapse of its roof, forming the 17-by-32-km oval depression of Long Valley caldera. The resulting ash-flow deposits, the Bishop Tuff, inundated 1,500 km^2 of the surrounding countyside and accumulated locally to thicknesses approaching 200 m on the Volcanic Tableland in upper Owens Valley, in Adobe Valley, and in the southwestern part of Mono Basin. Some ash flows spilled westward over the crest of the Sierra

NEW YORK
JUL 5 1989
STATE LIBRARY

best copy available

Nevada, filling the canyons of the Middle Fork of the San Joaquin River, although subsequent erosion has removed most of this intercanyon tuff. A large volume of the Bishop Tuff also ponded within the caldera, where drill holes (Smith and Rex, 1977; Gambill, 1981) have penetrated as much as 1,500 m of the Bishop Tuff buried beneath younger caldera fill. Plinian ash clouds associated with this climactic eruption drifted thousands of kilometers downwind, depositing an ash layer (informally designated the Bishop ash bed) recognized as far east as Kansas and Nebraska (Izett and others, 1970; Izett, 1982). Thinner Bishop ash deposits also are recognized in southwest California (Merriam and Bischoff, 1975) and in Pacific Ocean cores (Sarna-Wojcicki and others, 1987). Gilbert (1938) first mapped and described the Bishop Tuff and recognized Long Valley as its probable source. Hildreth and Mahood (1986) have shown, on the basis of the distribution of accidental inclusions in the deposits, that the initial Plinian eruptions vented in the south-central part of the caldera and that, with the transition to ash-flow eruptions, venting migrated around the ring-fracture zone, the final eruptions occurring along the northern caldera margin.

The Bishop Tuff is composed of porphyritic rhyolite ash and pumice lapilli and blocks containing 5 to 25 percent phenocryts of sanidine, quartz, plagioclase, biotite, augite, hypersthene, Fe-Ti oxides, and accessories, and it shows considerable variation in degree of welding and crystallization (Sheridan, 1965, 1970). Exposures in the Owens River Gorge show evidence of multiple cooling units (Sheridan, 1967), suggesting that the tuff was emplaced during at least two major eruptive pulses. Hildreth (1979), on the basis of detailed mineralogical and chemical studies, has shown that the tuff represents the stratigraphically inverted contents of the upper part of the Long Valley magma chamber and that it preserves chemical gradients developed in the chamber prior to eruption.

After collapse of the roof of the magma chamber, eruptions continued within the 2– to 3–km–deep caldera. Early postcaldera pyroclastic eruptions formed a thick sequence of intracaldera bedded tuffs that were followed by extrusion of thin, hot, fluid obsidian flows. Simultaneously, renewal of magma presssure at depth uplifted, arched, and faulted the central part of the caldera floor, forming a resurgent dome (Smith and Bailey, 1968) about 10 km in diameter, 500 m high, and having a northwest-trending medial graben 4 km wide. This uplift occurred within 100,000 years or less after caldera collapse, between about 0.7 and 0.6 Ma, as indicated by K–Ar ages of the contemporaneous obsidian flows (Bailey and others, 1976). The rhyolite tephra and obsidian flows erupted during this episode are informally referred to as *early rhyolite* and are typically aphyric to sparsely porphyritic, containing less than 5 percent crystals of plagioclase, hypersthene, biotite, and Fe–Ti oxides. They contrast strikingly with the preceding crystal-rich Bishop Tuff and appatently erupted at higher temperatures and probably from greater depth.

During rise of the resurgent dome and eruption of the early rhyolite, the caldera was filled by a large lake, Pleistocene Long Valley Lake (Mayo, 1934), terraces and strandlines of which are well preserved along the east wall of the caldera and locally on the flanks of the resurgent dome. Continued rise of the resurgent dome eventually raised the lake surface to the level of the low southeast caldera rim, where overflow subsequently cut the Owens River Gorge. Gradual downcutting of the gorge, accompanied by intermittent tectonic tilting of the southeast rim near the Sierran front, ultimately drained the lake between 100,000 and 50,000 years ago. (Lake Crowley is not a remnant of the Pleistocene lake but a modern reservoir impounded behind a dam contructed in 1940.) During the early life of the Pleistocene lake, glaciers flowing into it from the High Sierra generated debris-laden icebergs that drifted across the lake, depositing large erratics of Sierran granite on the flanks of the resurgent dome, which stood as an island, and on the lake terraces on the east caldera wall.

After an interiude of about 100,000 years of volcanic quiescence, crystal-rich rhyolite began erupting in the caldera moat peripheral to the resurgent dome. This porphyritic rhyolite, informally referred to as *moat rhyolite*, typically contains 15 to 20 percent phenocrysts of plagioclase, quartz, sanidine, biotite, hornblende, and Fe-Ti oxides. It formed thick, steep-sided domes and flows, indicating higher viscosity and lower temperature than the early rhyolite, and probably signaled the onset of cooling and crystallization of the magma chamber. Moat rhyolite erupted at about 200,000-yr intervals—at 0.5, 0.3, and 0.1 Ma in clockwise succession around the resurgent dome, in the north, southeast, and west sectors of the moat, respectively—with four to five domes or flows erupting during each episode. The moat rhyolite consists of two distinct chemical types, low- and high-silica types, which erupted alternately in postcaldera time. The low-silica type (hornblende-biotite rhyolite, units Qmr1, 2, 3) is the most voluminous and most frequently erupted and probably represents the dominant volume (Hildreth, 1981) of the magma chamber. In contrast, the high-silica type (units Qmrh and Qmrm) has chemical characteristics similar to those of the Bishop Tuff, which suggests late development of Bishop-type magma in the upper part of the postcaldera magma chamber (Bailey, 1984). The 0.1-Ma western moat rhyolites appear to be the youngest lavas so far derived from the Long Valley chamber. However, seismological studies (Hill, 1976; Steeples and Iyer, 1976; Ryall and Ryall, 1981; Sanders, 1984; Hill and others, 1985b; Leutgert and Mooney, 1985) suggest that a residual body of magma still underlies the resurgent dome at depths of 5 to 15 km or more, providing a potential source for future eruptions. For such a body of silicic magma to have remained partially liquid and been a source for continuing eruptions throughout the past 700,000 years, additional heat must have been repeatedly added, most likely by periodic injection of basalt from the mantle into the lower part of the silicic

chamber (Lachenbruch and others, 1976). Recent tectonic and apparent magmatic unrest within Long Valley caldera (Hill and others, 1985a) suggests that a brief episode of such magmatic injection may have occurred between 1980 and 1986 (Bailey, 1984).

Mono–Inyo Craters volcanic chain

The volcanism associated with Long Valley caldera, beginning about 3.6 Ma with precaldera trachybasalt-trachyandesite and quartz latite, continuing with the rhyolite of Glass Mountain and climactic eruptions of the Bishop Tuff, and concluding with the early- and moat-rhyolite extrusions as young as 0.1 Ma, constitutes a generally mafic-to-silicic sequence. The Mono-Inyo Craters volcanic chain, a 45-km north-trending zone of vents extending from Mammoth Mountain through the west moat of the caldera to Mono Lake, overlaps this Long Valley sequence both spatially and temporally. It also may be interpreted as a mafic-to-silicic sequence derived from younger chambers probably not directly related to Long Valley chamber, although in the area of overlap the interrelation is ambiguous. As here defined, the Mono–Inyo Craters volcanic chain includes postcaldera basaltic and quartz-latitic lavas (including the dacites and rhyodacites in and near Mono Lake) whose vents are localized along the same general north-south trend as the Mono-Inyo Craters rhyolites. Although the actual fissures and faults along which these vents are aligned are evident only sporadically along the chain, they clearly parallel a pervasive north- to northnortheast-trending fracture system in the pre-Tertiary Sierran rocks that is evident in aerial photographs, satellite imagery, and digital topographic maps of the Sierra Nevada.

The Mono-Inyo sequence began between 0.3 and 0.2 Ma with the eruption of postcaldera trachbasalt-trachyandesite from numerous vents in and near the west moat of the caldera, including those at Devils Postpile National Monument, southwest of the caldera. These mafic flows filled the west moat locally to depths of as much as 250 m and poured around the resurgent dome sending long tongues of lava into the northern and southern parts of the caldera moat. Younger mafic vents erupted successively farther north near June Lake (40-20 ka?) and at Black Point on Mono Lake (13.3 ka)—suggesting that the fissure system propagated northward. Intermittently during intervals in this mafic activity, quartz-latite domes and flows vented locally in the western part of the caldera, as well as farther north in and near Mono Lake. The greatest volumes, however, accumulated near the northwest and southwest caldera margins, where the north-south fissure system apparently intersected caldera ring fractures. On the southwest edge of the caldera, between about 200 and 50 ka, extrusion of at least 12 quartz-latite and associated low-silica sodic rhyolite domes and flows built Mammoth Mountain, an imposing cumulovolcano straddling the caldera rim.

Rhyolites began erupting to form the Mono-Inyo Craters about 40–38 ka—first at the Mono Craters and more recently at the Inyo Craters. The arcuate chain of 30 or more overlapping rhyolite domes, flows (coulees), and craters that forms the Mono Craters apparently erupted along the mylonitized border of a subcircular Cretaceous pluton centered on Aeolian Buttes (Kistler, 1966a; see maps A and D). Seismological studies (Achauer and others, 1986) suggest that their probable magmatic source is a subjacent north-south elongate chamber at 10–20 km depth; there appears to be no evidence for a larger deep-crustal chamber. The older Mono domes (40,000–3,000 yr B.P.) are moderately to sparsely porhyritic, whereas the younger ones (3,000–550 yr B.P.) are aphyric, suggesting a recent increase in eruptive temperature or decrease in depth of eruptive source, or both. Radiocarbon dating and dendrochronological studies indicate that the youngest eruptions in the chain occurred at the northern end, most likely between A.D. 1325 and 1365 but no later than A.D. 1368 (Sieh and Bursik, 1986).

The Inyo Craters chain forms a discontinuous 10-km-long line of rhyolite dome-flows and craters that include Wilson Butte, Obsidian Dome, Glass Creek dome, north and south Deadman Creek domes, as well as several other smaller unnamed domes and craters. Wilson Butte and north Deadman Creek dome are chemically and petrographically similar to the rhyolites of the Mono Craters and also are significantly older than the others, about 1,350 and 6,000 yr B.P., respectively (Miller, 1985). Although within the Inyo Craters chain, they are considered southern outliers of the rhyolite of Mono Craters. The younger Inyo domes—Obsidian Dome, Glass Creek dome, south Deadman Creek dome, two smaller unnamed domes, and their associated tephras—all erupted within a span of probably no more than a few years, or possibly months. Radiocarbon dating of the tephra deposits indicates an age of 650–550 yr B.P. (Miller, 1985), but stratigraphic and dendrochronological considerations indicate that they commenced erupting after cessation of the northern Mono Craters eruptions, possibly A.D. 1368 (Sieh and Bursik, 1986), and could be as young as A.D. 1472 (Miller, 1985). The initial pyroclastic eruptions from the three main younger vents blanketed the sourrounding terrain with thick deposits of pumice lapilli and ash; this explosive activity ceased from all three vents before extrusion of any of the lava domes (Miller, 1985). Drilling (Eichelberger and others, 1985) has confirmed that this activity was fed by a shallow rhyolite dike, as suggested by Miller (1985).

The younger Inyo domes consist of two contrasting, fluidly intermixed rhyolites: one light-colored, pumiceous, and very coarsely porphyritic, and the other more silicic, sparsely porphyritic obsidian. The two types superficially resemble the porphyritic moat rhyolite of Long Valley caldera and the aphyric to sparsely porphyritic obsidian of the Mono Craters respectively and one possible explanation for their origin is that they represent the mixing of magmas from the Long Valley and Mono Craters chambers along a connecting fissure (Bailey and others, 1976; Sampson, 1987; Sampson and Cameron, 1987).

best copy available

During or shortly after the pyroclastic phase of the Inyo vents and probably before extrusion of the domes, phreatic explosions blasted three craters on the summit and south flank of Deer Mountain, a 0.1-Ma moat-rhyolite dome south of Deadman Creek dome (Rinehart and Huber, 1965). Two of these craters now contain small lakes (the Inyo Crater Lakes). Radiocarbon dating and dendrochronological considerations limit the eruptions between A.D. 1340 and 1460 (Wood, 1977). Several additional phreatic explosion craters of similar age erupted about 1 km to the northwest along a northwest-trending fault zone. The three aligned craters on Deer Mountain probably formed by the interaction of ground water with the same rhyolite dike that fed the Inyo domes, but no juvenile rhyolite magma reached the surface during the explosions; only hot mud and blocks of older trachyandesite, rhyolite, glacial debris, and basement rock were ejected. Other phreatic explosion craters and deposits occur 5 km to the south on the north face of Mammoth Mountain, where they are aligned subparallel to the caldera wall, apparently on a subsidiary caldera ring fracture. The associated phreatic debris is overlain by Inyo Craters pumice-fall deposits, but a radiocarbon age of 500 ± 200 yr B.P. obtained from a stump buried in the debris suggests that the eruptions were approximately contemporanceous with those from the Inyo Craters.

Still younger eruptions occurred at the northern end of the Mono-Inyo chain. In Mono Lake, several small dacite cinder cones and flows form Negit Island and the northeast corner of Paoha Island; a partly submerged rhyolite flow forms a small group of islets northeast of Negit; on the east side of Paoha is a small hydrothermally altered rhyodacite dome and a cluster of partly submerged phreatic explosion craters; and on the northern third of Paoha is a craggy rhyolite dome or invasive flow protruding through disrupted lake sediments (Lajoie, 1968). Most of Paoha Island is composed of lake sediments that were uplifted from the lake bottom, probably by intrusion of this latter rhyolite dome. Uplift caused slumping and sliding of these water-saturated sediments away from the center of the island, producing a remarkable landside jumble, particularly west of the island where the lake bathymetry suggests that slides extend under water at least 2 km laterally. Based on detailed study of radiocarbon-dated shorelines around Mono Lake, Stine (1984) concludes that the oldest of the dacites is slightly younger than 2,000 yr B.P. and that the youngest postdates 220 yr B.P.; uplift of Paoha Island and extrusion of the associated rhyolite is younger still and may have occurred between A.D. 1720 and 1850, according to Stine.

The bathymetry of Mono Lake (Scholl and others, 1967; Los Angels Department of Water and Power, 1987) shows that the lake is deepest immediately east and south of Paoha Island, where it attains depths of 40–50 m. Four or five small subcircular depressions on the lake bottom adjacent to Paoha Island suggest a peripheral arc of subaqueous craters. In all, there are at least ten subaerial vents and possibly five subaqueous vents within Mono Lake.

The recency and greater frequency of eruptions along the Mono-Inyo Craters chain during the past 3,000 years, together with evidence for increased magma temperature and (or) shallower eruption depth (Bailey, 1982b) suggests that in spite of the recent (1980–1986) episode of unrest at Long Valley, future eruptions are more likely to occur along the Mono-Inyo chain than at Long Valley (Hill and others, 1985a). The potential hazards associated with possible future eruptions in the area are outlined by Miller and others (1982).

STRUCTURE

As noted above, Long Valley caldera lies at the northern end of the Owens Valley rift, between the east-facing Sierran escarpment and the west-facing White Mountain front. The pre-Tertiary basement in the area is principally Sierran batholithic rocks and associated metamorphic roof pendents (map A). In the eastern half of the area, the basement consists mainly of Triassic granodiorite, which locally encloses masses of Paleozoic metasedimentary rocks and is intruded by small stocks of Jurassic granite. In the western half of the area, the basement consists mainly of Paleozoic metasedimentary rocks of the Mount Morrison roof pendant and metavolcanic rocks of the Ritter Range roof pendant, which dip nearly vertically and trend generally northwest and are intruded by moderately large Cretaceous granitic plutons. The contact between these contrasting eastern and western basement assemblages passes through the middle of Long Valley caldera, approximately coincident with the Hilton Creek fault (map B), although control for this contact is provided by only two deep intracaldera drill holes aided by the aeromagnetic expression of the sedimentary roof pendant rocks (Kane and others, 1976).

Structural trends in the immediate vicinity of the caldera are dominated by the northwest-trending Sierran frontal fault zone (map B), which parallels the northwest elongation of the Sierran plutons and associated roof pendants. Within the caldera, this northwest regional trend is reflected by the northwest trend of the medial graben on the resurgent dome, as well as by alignment of many of the intracaldera vents (map D). East of the caldera, structural trends are more northerly, subparallel to the White Mountain front, which lies 15 km east of the map area. Formation of the Sierran escarpment and most other Basin and Range escarpments in the area mainly postdates eruption of the precaldera lavas (3.6 Ma), and most of the bounding faults continue to be active, as attested by abundant and occasionally strong earthquakes along them (van Wormer and Ryall, 1980; Ryall and Ryall, 1980; Sherburne, 1980).

Major structual elements east of the caldera include the Benton Range, Casa Diablo Mountain, and Black Mountain—east-tilted fault-blocks bounded on the west by the Benton Range-Casa Diablo Mountain fault zone and the Black Mountain fault, respectively (map B).

These faults, and their many subsidiary splinters, displace Tertiary precaldera trachybasalt flows locally as much as 500 m; they also displace the rhyolite tuff fans of Glass Mountain and the Bishop Tuff by lesser amounts.

Major Sierran frontal faults within the map area include the Round Valley, Hilton Creek, Hartley Springs, Lee Vining, Fern Lake, and Silver Lake faults, all with displacements ranging from 500 to greater than 1,000 m down to the east. The Hilton Creek and Round Valley faults merge south of the map area to form the main west boundary of the Owens Valley rift. Northward, the Hilton Creek fault is defined by an anastomosing system of minor intracaldera faults that project into the Alpers Canyon fault and the Bald Mountain graben, both of which die out north of the caldera—a relation that suggests the precaldera Hilton Creek fault decreased in throw northward and terminated within or just north of the present site of the caldera. Crossing and diverging northnortheastward from the intracaldera continuation of the Hilton Creek fault is a narrow, segmented, grabenlike structure, starting on the northwest sector of the resurgent dome and contunuing north of the caldera as the Sage Hen Peak fault and associated graben. This structure merges with the Baxter Spring and Indian Spring faults, which define a northnorthwest-tending graben that narrows northward and continues as a single prominent west-downward normal fault east of Mono Lake. This complex but more or less continuous zone transecting the east-central part of the caldera consists essentially of two major northnorthwest-trending normal faults, the Hilton Creek in the south and the Indian Springs in the north, downthrown east and west respectively, joined by a northnortheast-trending graben, the Sage Hen Peak fault and graben.

Transecting the western part of the caldera is the Hartley Springs fault, which lies on strike with the Lee Vining fault to the north but is separated from it by an apparent structural gap through Pumice Valley. The Hartley Springs fault projects southward through the west moat of the caldera toward the Laurel-Convict fault, but the latter is a Paleozoic of Mesozoic structure within the Mount Morrison roof pendant with no apparent Cenozoic displacement. Continuation of the Hartley Springs fault through the caldera is suggested by the alignment of west moat rhyolite domes (maps B and D) and by a slight oversteepening of west moat trachybasalt flows along a coincident line, suggesting that they overflowed a preexisting east-downward fault scarp. The decline in displacement along the Lee Vining-Hartley Springs zone from greater than 1,000 m near Lee Vining to 600 m near the caldera suggests that the fault dies out southward, just as the Hilton Creek fault does northward to the east. This en echelon relation of the Hartley Springs and Hilton Creek faults suggests the existence of a northerly inclined ramp between the two faults in precaldera time.

The Silver Lake-Fern Lake fault zone intersects the west wall of the caldera, and the alignment of several postcaldera trachybasalt-trachyandesite vents in the west and southwest moat (map D) suggests that its intracaldera continuation crosses the inferred intracaldera segment of the Hartley Springs fault and arcs into the south moat, where it possibly coincides with the caldera ring fault. This inference suggests that in precaldera time, in the present site of the south moat, the eastward extention of the Fern Lake fault formed a north-downward east-west scarp across the north-sloping ramp between the Hartley Springs and Hilton Creek faults.

Transecting these converging structures in the west moat is a young north-trending, right-stepping, en echelon graben related to the Mono-Inyo Craters fissure system. This structure narrows and continues south of the caldera as a zone of minor faults and fissues in the granite of the High Sierra, along which is located Fish Creek (Iva Bell) Hot Springs, near the south edge of the Devils Postpile quadrangle, and possibly also Mono Hot Springs, about 15 km farther south in the Kaiser Peak quadrangle.

Both the Hartley Springs and Hilton Creek faults terminate as major escarpments at the north and south caldera boundaries, respectively, and continue within the caldera only as minor, discontinuous, en echelon splinters. This marked difference in displacement outside and inside the caldera suggests that in postcaldera time the cauldron block was structurally isolated by the ring fault and that the residual subcauldron magma chamber hydraulically dampened or absorbed intracaldera fault movements; possibly only recently has the chamber roof thickened sufficiently, as a result of cooling and crystallization, to behave rigidly and transmit tectonic stresses across the caldera floor. However, incomplete knowledge of the relative amounts of pre-, syn-, and post-caldera throw on these Sierran faults prevents full development of this hypothesis. A similar difference in displacement is evident farther north where Pumice Valley embays the Sierran front west of the Mono Craters; here both the Lee Vining and Hartley Springs faults terminate as major escarpments at the north and south edges of Pumice Valley. However, there appears to be no large shallow magma body beneath Pumice Valley to provide a similar explanation. An alternative explanation, which may apply to Long Valley as well, is offered by Burisk (Sieh and Bursick, 1986), whereby the space generated by extension along the Sierran front west of the Mono Craters is occupied by dike intrusion beneath the craters, or in the case of Long Valley, by periodic refilling of the magma chamber.

Development of the Long Valley magma chamber and location of the caldera were undoubtedly controlled by structural factors. The caldera clearly formed along a segmant of the Sierran frontal fault where left-stepping en echelon offset is unusually well developed and where the northwest-trending Sierran front diverges from the more northerly trend of the White Mountains front. Both of these relations suggest a region of localized extention that would facilitate rise of mafic magma from depth and where opening of fractures would promote magma stoping, fusion, and

best copy available

fractionation and provide space for the more silicic melts thus generated.

The main Long Valley caldera ring fault, presumably a subcircular or elliptical fault or fault zone bounding the subsided cauldron block, is not exposed, but its general location, coincident with the annular caldera moat, is suggested by the distribution of the moat rhyolite vents peripheral to the resurgent dome. The main ring fault appears to lie well within the present topographic boundary, which was enlarged by landslides, avalanches, and rapid erosion of the initially steep, fault-bounded walls during or shortly after caldera collapse. In the moat, on the east side of Long Valley proper, the position of the main ring fault is suggested by an arc of discontinuous minor faults—the Long Valley fault zone (map B). Although not a prominent escarpment, and possibly in part the result of differential compaction of thick intracaldera sediments, this zone coincides with the steep arcuate gravity gradient (Kane and others, 1976) that is inferred to define the main structural boundary in the subsurface. East of this apparent main fault zone are two subsidiary ring faults, the Watterson Canyon fault, which approximately coincides with the present topographic rim of the caldera, and the Clover Patch fault, which is considerably east of the topographic rim. These subsidiary arcuate faults, which are unique to the eastern side of the caldera, are possibly a consequence of the laccolithic form of the Long Valley magma chamber postulated by Lachenbruch and others (1976), wherein evacuation of the shallow, eastern part of the chamber generated lateral rotational sliding of megaslide blocks rather than vertical collapse during caldera subsidence.

Evidence for postsubsidence modification and enlargement of the caldera is best recorded along this eastern margin. Here, the topographic rim is 3-4 km east of the apparent main ring fault. Immediately south of Glass Mountain the rim coincides with the trace of the Watterson Canyon fault; farther south the rim diverges eastward as much as 2 km from the fault trace and forms the edge of the Volcanic Tableland east of Lake Crowley. In this area the fault trace is partly buried by postcaldera sediments, which show that thick rhyolite tuffs of Glass Mountain (unit Qgt), initially forming the fault-bounded rim, were rapidly eroded and redeposited as coarse alluvial fans (unit Qof). These fan deposits in turn were reworked by the caldera lake into tiers of terrace gravels (unit Qtg; see cross section C-C'). The relations indicate that the topographic rim, particularly east of Lake Crowley, has migrated eastward of both the Long Valley and Watterson Canyon faults and is in large part a fault line scarp. In the western, steep-walled part of the caldera, indirect evidence for landslide enlargement of the walls is found in collapse megabreccias (Lipman, 1976) encountered within the Bishop tuff in at least one intracaldera drill hole (No. 5, table 1).

Table 1.—*Principal drill holes in Long Valley-Mono Craters area*

Map No.	Date	Number	Operator/Location	Elevation (m)	Depth (m)	Temperature Max./Bottom (degrees C)		Reference
1	(1971)	PRC 4397.1/1	Geothermal Resources International south shore Mono Lake, T. 1 N., R. 27 E., sec. 20	1954	1253	46	46	Axtell (1972)
2	(1971)	PRC 4372.1/23–1	Getty Oil north shore Mono Lake T. 2 N., R. 26 E., sec. 23	1957	743	58	58	Axtell (1972)
3	(1976)	LV 66–29	Republic Geothermal central Long Valley T. 3 S., R. 29 E., sec. 29	2128	2109	72	72	Smith and Rex (1977)
4	(1979)	CP–1	Union Oil of California east flank, resurgent dome T. 3 S., R. 28 E., sec. 15	2226	1846	142	103	Gambill (1981)
5	(1979)	M–1	Union Oil of California Casa Diablo Hot Springs T. 3 S., R. 28 E., sec. 32	2226	1605	146	146	Gambill (1981)
6	(1982)	82–6	Union Oil of California southeast of Mono Craters T. 1 S., R. 27 E., sec. 33	2341	899	28	28	R.F. Dondanville, written commun., 1987
7	(1982)	PLV–1	Phillips Petroleum west moat of caldera T. 3 S., R. 27 E., sec. 22	2585	715	110	110	Benoit (1984)
8	(1982)	PLV–2	Phillips Petroleum west moat of caldera T. 3 S., R. 27 E., sec. 3	2347	636	68	68	Benoit (1984)
9	(1985)	IDFU 44–16	Unocal Geothermal west moat of caldera T. 3 S., R. 27 E., sec. 16	2451	1796	215	180	Suemnicht, 1987
10	(1986)	SR (RDO–8)	Lawrence Berkeley Laboratory Shady Rest, Mammoth Lakes T. 3 S., R. 27 E., sec. 25	2374	715	202	202	Wollenberg and others (1987)

Postsubsidence intracaldera volcanic vents show a strong preference for alignment along Sierran frontal trends (map D). Most early rhyolite vents are localized along northwest-trending faults bounding the medial graben. Although the moat rhyolites are arranged peripheral to the resurgent dome, individual vents particularly those north and west of the resurgent dome, are aligned along intracaldera projections of the Hilton Creek and Hartley Springs faults, respectively. Similarly, several postcaldera trachybasalt-trachyandesite vents are aligned along the intracaldera projection of the Fern Lake fault. In contast, vents for the Mono-Inyo Craters volcanic chain trend generally north-south, although those at the soutern end near Pumice Butte form a broad zone that trends more northnortheasterly. The rhyolite vents of the Mono–Inyo Craters chain form a generally linear northerly trend, but locally they appear to consist of northnortheast-trending en echelon segments (Fink, 1985). Vents of the central segment of the Mono Craters arc appear to be aligned on such a trend, but the curvature shown by the vents at the northern and southern extremities, together with the location of the granodiorite-metasedimentary rock contact beneath the craters, as exposed in the Mono Craters tunnel (map A), reflects the structural influence of the mylonitized border of the Aeolian Buttes pluton (Kistler, 1966b).

Precaldera volcanic vents do not show any strong preferential alignment (map C). They tend to be concentrated around the northern margin of the caldera, but this probably is more apparent than real, as over much of the area precaldera rocks are buried by the Bishop Tuff and younger tephra, and within the caldera they are downfaulted and buried by caldera fill. It is evident, however, that quartz latite vents are concentrated near the caldera, whereas basaltic vents are more widely distributed. Both north and northeast vent alignments are evident locally, but neither trend is dominant or traceable for far due to younger cover. Few vents, either basaltic or quartz latitic, occur immediately south of the caldera, but to the southwest, regional mapping (Moore and Dodge, 1980; Bailey, 1982a) shows that 20 or more basaltic vents are scattered over a large area in the High Sierra. The precaldera rhyolites of Glass Mountain form an arcuate ridge along the northeast caldera rim, but the actual vents do not have a demonstrably arcuate pattern clearly contolled by incipient caldera ring fractures; as previously suggested (Bailey and others, 1976). Many additional rhyolite vents probably are buried within the caldera.

HOT SPRINGS, FUMAROLES, AND THERMAL ACTIVITY

Long Valley caldera is recognized as a region of high heat flow (Lachenbruch and others, 1976), and hot springs, fumaroles, and active hydrothermal alteration are prevalent in many parts of the caldera, particularly in the southeast moat and adjacent flanks of the resurgent dome. Notable concentrations of fumaroles and (or) hot springs occur at Casa Diablo Hot Springs, Fumarole Valley, Hot Creek Springs, Little Hot Creek Springs, and southeast of Big Alkali and Little Alkali Lakes. Fine-grained black pyrite (FeS_2) and travertine ($CaCO_3$) are being actively deposited in many of these springs, and native sulfur is a common sublimate around fumaroles. Siliceous sinter also occurs in some of these areas, but it is the result of former higher temperature hot-spring activity and is not forming at the present time. (An isolated occurrence of siliceous sinter is reported by Krauskopf and Bateman, (1977) in the faulted Tertiary basalt terrane in the far northeast corner of the map area, but its age and relation to possible thermal sources is not known.) The coarse, silica-cemented, deltaic sediments (unit Qsc) that flank the east and south sides of the resurgent dome are pervasively penetrated by fossil fumarolic pipes, suggesting an unusually intense episode of thermal activity during deposition of this unit about 300,000 yr ago. Quite possibly, eruption of the southeast moat rhyolites at about this time was accompanied by renewed uplift of the resurgent dome and by intensified hydrothermal activity, thereby causing the shedding of coarser detritus off the resurgent dome into the caldera lake and concomitant widespread silica-cementation of the deltaic sediments thus formed. Pervasive hydrothermal alteration of lake beds beneath the deltaic sediments is evident near many faults on the resurgent dome. This alteration is predominantly argillic, and near Little Antelope Valley commercial grade kaolin has been mined intermittently since the 1950's (Cleveland, 1962).

The prevalence of thermal activity in Long Valley has encouraged geothermal exploration, and several commercial and scientific exploratory holes have been drilled to depths ranging from 100 to 2,000 m. (The location of drill holes deeper than 500 m for which data are available are shown on the geologic map and cross sections and are keyed by number to supplementary data and references presented in table 1.) Chemical geothermometry of hot spring and drill-hole waters suggests that deep reservoir temperatures range from 200 to 280 °C, but the maximum temperature attained in drill holes is 220 °C (Suemnicht, 1987). Drilling to date (1987) indicates that thermal waters are confined to relatively shallow aquifers, mainly within the intracaldera Bishop Tuff, at depths above 1,500 m in the west moat and above 700 m in the south moat, but neither the location of the deep thermal source nor where or how the aquifers are fed is known. Presumably residual magma beneath the resurgent dome is the principal heat source, but higher temperatures at greater depths in the western part the of the caldera suggest a possible westerly source (Sorey, 1978, 1985; Blackwell, 1985).

Surface thermal acativity associated with the Mono–Inyo Craters volcanic chain is restricted to the very southern and northern extremities of the chain. In the south, active fumaroles occur on the north and south slopes of Mammoth Mountain, and hot springs occur to the west at Reds Meadow Hot Springs and to the south in Fish Creek at Iva Bell Hot Springs. In the vicinity of Mono Lake, fumaroles and hot springs occur on Paoha

best copy available

Island, and hot springs occur northeast of Black Point cinder cone. Although surface thermal features are confined to the extremeties of the chain, elevated temperatures do occur in the subsurface along it. During the course of constructing the Mono Craters tunnel (map A), warm water (35 °C) and copious amounts of carbon dioxide were encountered when passing through feeder dikes (Gresswell, 1940), and a bottom-hole temperature of 80 °C was attained in the Obsidian Dome conduit drill hole in the Inyo Craters chain (Eichelberger, and others, 1985).

GLACIATION

The oldest currently recognized glacial deposit in the Long Valley area is the McGee Till (unit Qmg) (Blackwelder, 1931), exposed on the south rim of the caldera on McGee Mountain. It overlies a scoriaceous basaltic vent complex having a radiometric age of 2.6 Ma (Dalrymple, 1963), and although generally considered Pleistocene in age, it could be older. The 3-Ma here-abandoned Deadman Pass Till (Curry, 1966) is now recognized as a product of erosion and weathering of pyroclastic deposits unusually rich in accidental granitic and metamorphic clasts (R.A. Bailey, N.K.Huber, and R.R. Curry, unpub. data, 1987), and it is here mapped as part of the quartz latite of San Joaquin Ridge (unit Tsjt).

During Sherwin glaciation (Blackwelder, 1931), thick, now much eroded, lobate masses of till were deposited at many places along the Sierran front, particularly at the mouth of Rock Creek (Putnam, 1960; Sharp, 1968) and south of Lee Vining Creek (Kistler, 1966b). Scattered remnants of the Sherwin till or related outwash deposits also occur beneath the Bishop Tuff at Aeolian Buttes (Putnam, 1949), between precaldera basalt and the Bishop Tuff near Crestview, and between older Quaternary gravels (unit Qog) and Bishop Tuff near Alpers Canyon on the northwest rim of the caldera. Sporadic granitic blocks, also possible remnants of Sherwin outwash, rest on precaldera basalt east of Bald Mountain in upper McLaughlin Creek drainage. Sherwin outwash, clearly related to the Sherwin Till at the mouth of Rock Creek, also crops out between Tertiary basalt and the Bishop Tuff in the walls of Owens River Gorge (Putnam, 1960; Sharp, 1968). The age of the Sherwin Till is not known precisely, but on the basis of depth of weathering of the till beneath the Bishop Tuff, Sharp (1968) estimates that it may predate the tuff by a few ten-thousand years but not by as much as 100,000 years.

Fragmentary evidence suggest that glaciers were active in the Sierra Nevada at the time of eruption of the Bishop Tuff and for some time after formation of Long Valley caldera. Whether this glaciation was a continuation of the Sherwin glaciation or a distinctly younger stage is uncertain. Evidence for glaciation coeval with emplacement of the Bishop Tuff is displayed at Little Round Valley, just south of Lake Crowley, where a remarkable display of fossil fumarolic pipes and other unusual depositional features in the tuff suggest the proximity of a ice-tuff contact (R.A. Bailey, unpub. data, 1986). Clear evidence for active glaciation during early postcaldera time is provided by the ice-rafted erratics dispersed on early intracaldera lake terraces.

Glacial deposits of Wisconsin age in the Long Valley-Mono Basin area have been studied for many decades, but their subdivision and correlation remain problematic and are to some degree controversial. The glacial subdivisions shown on this map are based on casual observations made over several years, plus two weeks of intensive comparative examination of most of the moraines in the map area, qualitatively applying the semiquantitative critera of Sharp (1969). In most drainages, the subdivisions shown are in general agreement with recent studies; where they differ, or where subdivisions have not been previously made, they are commonly queried (for example, Qmb? in Laurel, Sherwin, and Convict Creeks), and they require verification by more detailed study. In the Long Valley area, from Rock Creek northwest to Mammoth Creek, two distinct morainal lobes can be readily distinguished in most of the major Sierran valleys; in general agreement with most previous work, the older and younger lobes are mapped as deposits of Tahoe and Tioga glaciations, respectively. In Mammoth Creek, where Wisconsin age glaciers spread out as broad piedmont lobes, the youngest moraine overlies a trachybasalt flow with a radiometric age of 64 ka (Bailey and others, 1976) and is mapped as Tioga in age. The Casa Diablo Till (Curry, 1971) lies between trachybasalt and trachanydesite flows equivocally dated at 64 and 126 ka, respectively; it may be Tahoe or Mono Basin in age. In the Mono Basin area, multiple Wisconsin - age glaciations are more readily distinguishable than in Long Valley, three or four distinct pairs of lateral moraines being recognizable in most of the major valleys. These are mapped as deposits of Mono Basin, Tahoe, Tenaya, and Tioga glaciations, in general agreement with mapping by Sharp and Birman (1963) and Gillespie 1982).

REFERENCES CITED

Achauer, Ulrich, Greene, Lizbeth, Evans, J.R., and Iyer, H.M., 1986, Nature of the magma chamber underling the Mono Craters area, eastern California, as determined from telesismic traveltime residuals: Journal of Geohysical Research, v. 91, no. B14, p. 13,873–13,891.

Axtell, L.H., '1972, Mono Lake geothermal wells abandoned: California Geology, v. 25, no. 3, p. 66–67.

Bailey, R.A., 1982a, Geothermal-resource evaluation of the Minarets Wilderness and adjacent areas, Madera and Mono Counties, California: U.S. Geological Survey Bulletin 1516-C, p. 73-89.

———1982b, Other potential eruption centers in California: Long Valley-Mono Lake, Coso, and Clear Lake volcanic fields: California Division of Mines and Geology Special Publication 63, p. 17–28.

———1984, Chemical evolution and current state of the Long Valley magma chamber, *in* Hill, D.P.,

Bailey, R.A., and Ryall, A.S., eds., Proceedings of Workshop XIX: Active tectonic and magmatic processes beneath Long Valley caldera, eastern California: U.S. Geological Survey Open-File Report 84–939, v. 1, p. 24–40.

Bailey, R.A., Dalrymple, G.B., and Lanphere, M.A., 1976, Volcanism, structure, and geochronology of Long Valley caldera, Mono County, California: Journal of Geophysical Research, v. 81, no. 5, p. 725–744.

Benoit, W.R., 1984, Initial results from drillholes PLV-1 and PLV-2 in the western moat of the Long Valley caldera: Geothermal Resources Council Transactions, v. 8, p. 397–402.

Birman, J.H., 1964, Glacial geology across the crest of the Sierra Nevada, California: Geological Society of America Special Paper 75, 80 p.

Blackwelder, Eliot, 1931, Pleistocene glaciation in the Sierra Nevada and Basin Ranges: Geological Society of America Bulletin, v. 42, p. 865–922.

Blackwell, D.D., 1985, A transient model for the geothermal system of Long Valley caldera, California: Journal of Geophysical Research, v. 90, no. B13, p. 11,229–11,241.

Burke, R.M., and Birkeland, P.W., 1979, Reevaluation of multiparameter relative dating techniques and their application to the glacial sequence along the eastern escarpment of the Sierra Nevada, California: Quaternary Research, v. 11, p. 21–51.

Chaudet, R.C., 1986, The petrology and geochemistry of precaldera magmas, Long Valley caldera, eastern California: Blacksburg, Virginia, Virginia Polytechnical Institute, M.Sc. thesis, 151 p.

Chesterman, C.W., and Gary, C.H., Jr., 1975, Geology of the Bodie quadrangle, Mono County, California: California Division of Mines and Geology, map sheet 21, scale 1:48,000.

Cleveland, G.B., 1962, Geology of the Little Antelope Valley clay deposits, Mono County, California: California Division of Mines and Geology Special Report 72, p. 1–28.

Curry, R.R., 1966, Glaciation about 3,000,000 years ago in the Sierra Nevada, California: Science, v. 154, p. 770–771.

———1971, Glacial and Pleistocene history of the Mammoth Lakes Sierra, California: a geologic guidebook: Missoula, Montana, University of Montana Geological Series Publication, no. 11, 49 p.

Dalrymple, G.B., 1963, Potassium-argon dates of some Cenozoic volcanic rocks of the Sierra Nevada, California: Geological Society of America Bulletin, v. 74, p. 379–390.

———1972, Potassium-argon dating of geomagnetic reversals and North American glaciations, *in* Bishop, W.W., and Miller, J.A., eds., Calibration of Hominoid Evolution: Scottish Academic Press, p. 101–134.

Eichelberger, J.C., Lysne, P.C., Miller, C.D., and Younker, L.W., 1985, Research drilling at Inyo domes, 1984 results: EOS, American Geophysical Union Transactions, v. 66, no. 17, p. 186–187.

Fink, J.H., 1985, Geometry of the silicic dikes beneath the Inyo domes, California: Journal of Geophysical Research, v. 90, no. B13, p. 11,127–11,133.

Fiske, R.S., and Tobisch, O.T., 1978, Paleogeographic significance of volcanic rocks of the Ritter Range pendant, central Sierra Nevada, *in* Howell, D.G., and McDougall, K.A., eds., Mesozoic Paleogeography of the Western United States: Society of Economic Paleontologists and Mineralogists, Pacific Section, Los Angeles, California, p. 209–222.

Gambill, D.T., 1981, Preliminary hot dry rock geothermal evaluation of Long Valley caldera, California: Los Alamos National Laboratory Report, LA 8710-HDR, 22 p.

Gilbert, C.M., 1938, Welded tuff in eastern California: Geological Society of America Bulletin, v. 49, p. 1829–1862.

Gilbert, C.M., Christiansen, M.N., Al Rawi, Yehya, and Lajoie, K.R., 1968, Structural and volcanic history of Mono Basin, California-Nevada: Geological Society of America Memoir 116, p. 275–329.

Gillespie, A.R., 1982, Quaternary glaciation and tectonism in the southeastern Sierra Nevada, Inyo County, California: Pasadena, Calif., California Intstitute of Technology, Ph.D. dissertation, 695 p.

Gresswell, W.K., 1940, Short report on the geological information encountered in driving the Mono Craters tunnel: California Journal of Mines, v. 36, p. 199–204.

Hildreth, Wes, 1979, The Bishop Tuff: Evidence for the origin of compositional zoning in silicic magma chambers: Geological Society of America Special Paper 180, p. 43–75.

———1981, Gradients in silicic magma chambers: implications for lithospheric magmatism: Journal of Geophysical Research, v. 86, no. B11, p. 10,153–10,192.

Hildreth, E.W., and Mahood, G.A., 1986, Ring-fracture eruption of the Bishop Tuff: Geological Society of America Bulletin, v. 97, p. 396–403.

Hill, D.P., 1976, Structure of Long Valley caldera, California, from a seismic reflection experiment: Journal of Geophysical Research, v. 81, no. 5, p. 745-753.

Hill, D.P., Bailey, R.A., and Ryall, A.S., 1985a, Active tectonic and magmatic processes beneath Long Valley caldera, eastern California: an overview: Journal of Geophysical Research, v. 90, no. B13, p. 11,111–11,120.

Hill, D.P., Kissling, E., Luegert, J.H., and Kradolfer, U., 1985b, Constraints on the upper crustral structure of the Long Valley-Mono Craters volcanic complex, eastern California, from seismic refraction measurements: Journal of Geophysical Research, v. 90, no. B13, p. 11,135–11,150.

Huber, N.K., 1981, Amount and timing of Late Cenozoic uplift and tilt of the central Sierra Nevada, California—evidence from the upper San Joaquin River basin: U.S. Geological Survey Professional Paper 1197, 28 p.

Huber, N.K., and Rinehart, C.D., 1965, Geologic map of the Devils Postpile quadrangle, sierra Nevada,

best copy available

California: U.S. Geological Survey Geologic Quadangle Map GQ–437, scale 1:62,500.
———1967, Cenozoic volcanic rocks of the Devils Postpile quadrangle, eastern Sierra Nevada, California: U.S. Geological Survey Professional Paper 554–D, p. D1–D21.
Izett, G.A., 1982, The Bishop ash bed and some older compositionally similar ash beds in California, Nevada and Utah: U.S. Geological Survey Open-File Report 82–582, 47 p.
Izett, G.A., Wilcox, R.E., Powers, H.A., and Desborough, G.A., 1970, the Bishop Ash Bed, a Pleistocene marker bed in the Western United States: Quaternary Research, v. 1, p. 121–132.
Kane, M.F., Mabey, D.R., and Brace, Rosa-Lee, 1976, A gravity and magnetic investigation of the Long Valley caldera, Mono County, California: Journal of Geohpysical Research, v. 81, 5, p. 754–768.
Kistler, R.W., 1966a, Structure and metamorphism in the Mono Craters quadrangle, Sierra Nevada, California: U.S. Geological Survey Bulletin 1221–E, p. E1–E53.
———1966b, Geologic map of the Mono Craters quadrangle, Mono and tuolumne Counties, California: U.S. Geological Survey Geologic Quadrangle Map GQ–462, scale 1:62,000.
Kistler, R.W., and Nokleberg, W.J., 1979, Carboniferous rocks of the eastern Sierra Nevada, *in* Saul, R.B., Bowen, O.E., Stevens, C.H., Dunne, G.C., Randall, R.G., Kistler, R.W., Nokleberg, W.J., D'Allura, J.A., Moores, E.M., Watkins, Rodney, Baldwin, E.M., Gilmour, E.H, and Danner, W.R., The Mississippian and Pennsylvanian (Carboniferous) Systems in the United States—California, Oregon, and Washington: U.S. Geological Survey Professional Paper 1110-CC, p. CC21–CC26.
Kistler, R.W., and Swanson, S.E., 1981, Petrology and geochronology of metamorphosed volcanic rocks and a Middle Cretaceous volcanic neck in the east-central Sierra Nevada, California: Journal of Geophysical Research, v. 86, no. Bll, p. 10,489–10,501.
Krauskopf, K.B., and Bateman, P.C., 1977, Geologic map of the Glass Mountain quadrangle, Mono County, California, and Mineral County, Nevada: U.S. Geological Survey Geologic Quadrangle Map GQ–1099, scale 1:62,500.
Lachenbruch, A.H., Sass, J.H., Munroe, J.R., and Moses, T.H., 1976, Geothermal setting and simple heat conduction models for Long Valley caldera: Journal of Geophysical Research v. 81, no. 5, p. 769–784.
Lajoie, K.R., 1968, Late Quaternary stratigraphy and geologic history of Mono Basin, Eastern California: Berkeley, Calif., University of Californa, Ph.D. dissertation, 271 p.
———1969, Late Quaternary stratigraphy and geologic history of Mono Basin, eastern California: Dissertation Abstracts, v. 30, p. 1202.
Lipman, P.W., 1976, Caldera-collapse breccias in the western San Juan Mountains, Colorado: Geological Society of America Bulletin, v. 87, p. 1397–1410.
Los Angeles Department of Water and Power, 1986, A bathymetric and geologic survey of Mono Lake, California: Report No. 15, 38 p., 9 plates.
Luetgert, J.H., and Mooney, W.D., 1985 Earthquake profiles from the Mammoth Lakes, California, earthquake swarm, January 1983: Implications for deep crustal structure: Seismological Society of America Bulletin, v. 75, p. 211–221.
Mankinen, E.A., Gromme, C.S., Dalrymple, G.B., Lanphere, M.A., and Bailey, R.A., 1986, Paleomagnetism and K–Ar ages of volcanic rocks from Long Valley caldera, California: Journal of Geophysical Research, v. 91, no. B1, p. 633–652.
Mayo, E.B.,1934, The Pleistocene Long Valley Lake in eastern California: Science, v. 80, p. 95–96.
Merriam, Richard, and Bischoff, J.L., 1975, Bishop Ash: a widespread volcanic ash extended to southern California: Journal of Sedimentary Petrology, v. 45, p. 207–211.
Metz, J.M, and Mahood, G.A., 1985, Precursors to the Bishop Tuff eruption: Glass Mountain, Long Valley, California: Journal of Geophysical Research, v. 90, no. B13, p. 11,121–11,126.
Miller, C.D., 1985, Holocene eruptions at the Inyo volcanic chain, California: implications for possible eruptions in Long Valley: Geology v. 13, p. 14–17.
Miller, C.D., Mullineaux, D.R., Crandell, D.R., and Bailey, R.A., 1982, Potential hazards from future volcanic eruptions in the Long Valley-Mono Lake area, east-central California and Nevada—a preliminary assessment: U.S. Geological Survey Circular 877, 10 p.
Moore, J.G., and Dodge, F.C.W., 1980, Cenozoic volcanic rocks of the southern Sierra Nevada, California: I—Geology and petrology: Summary: Geological Society of America Bulletin, v. 91, p. 515–518.
Putnam, W.C., 1949, Quaternary geology of the June Lake disrtict, California: Geological Society of America Bulletin, v. 60, p. 1281–1302.
———1960, Origin of Rock Creek and Owens Rivers Gorges, Mono County, California: University of California Publications in Geology, v. 34, p. 221–280.
Rinehart, C.D., and Huber, N.K., 1965, The Inyo Crater Lakes—a blast in past: California Division of Mines and Geology, Mineral Information Service, v. 18, no. 9, p. 169–172.
Rinehart, C.D., and Ross, D.C., 1957, Geology of the Casa Diablo Mountain quadrangle, California: U.S. Geological Survey Geologic Quadrangle Map GQ–99, scale 1:62,500.
———1964, Geology and mineral deposits of the Mount Morrison quadrangle, Sierra Nevada, California, *with a section on* A gravity study of Long Valley, by L.C. Pakiser: U.S. Geological Survey Professional Paper 385, 106 p.
Ryall, A.S., and Ryall, F.D., 1980, Anomalous patterns of earthquake occurrence in the Sierra Nevada-Great Basin boundary zone, *in* Evernden, J.F., ed., Proceedings of Conference X: Earthquake hazards

along the Wasatch and Sierra Nevada frontal fault zones: U.S. Geological Survey Open-File Report 80–801, p. 348–358.

Ryall, F.D., and Ryall, A.S., 1981, Attenuation of P and S waves in a magma chamber in Long Valley caldera, California: Geophysical Research Letters, v. 8, p. 557–560.

Sampson, D.E., 1987, Textural heterogeneities and vent area structures in the 600-year-old lavas of the Inyo volcanic chain, eastern California: Geological Society of America Special Paper 212, p. 89–101.

Sampson, D.E., and Cameron, K.L., 1987, The geochemistry of the Inyo volcanic chain: multiple magma systems in the Long Valley region, eastern California: Journal of Geophysical Research, v. 92, no. B10, p. 10, 403–10,421.

Sanders, C.O., 1984, Location and configuration of magma bodies beneath Long Valley, California, determined from anomalous earthquake signals: Journal of Geophysical Research, v. 89, no. B10, p. 8287–8302.

Sarna-Wojcicki, A.M., Morrison, S.D., Meyer, C.E., and Hillhouse, J.W., 1987, Correlation of upper Cenozoic tephra layers between sediments of Western United States and the East Pacific Ocean, and comparison with biostratigrphic and magnetostratigraphic age data: Geological Society of America Bulletin, v. 98, p. 207–233.

Scholl, D.W., von Huene, Roland, St.-Amand, Pierre, and Ridlon, J.B., 1967, Age and origin of topography beneath Mono Lake, a remnant Pleistocene lake, California: Geological Society of America Bulletin, v. 78, p. 583–600.

Sharp, R.P., 1968, Sherwin Till-Bishop tuff geological relationships, Sierra Nevada, California: Geological Society of America Bulletin, v. 79, p. 351–364.

———1969, Semiquantitative differentiation of glacial moraines near Convict Lake, Sierra Nevada, California: Journal of Geology, v. 77, p. 68–91.

Sharp, R.P., and Birman, J.H., 1963, Additions to classical sequence of Pleistocene glaciations, Sierra Nevada, California: Geological Society of America Bulletin, v. 74, p. 1079–1086.

Sherburne, R.W., editor, 1980, Mammoth Lakes, California, earthquakes of May 1980: California Division of Mines and Geology Special Report 150, 141 p.

Sheridan, M.F., 1965, The mineralogy and petrology of the Bishop Tuff: Stanford, Calif., Stanford University, Ph.D. dissertation, 165 p.

———1967, Double cooling-unit nature of the Bishop Tuff in Owens River Gorge, California (abstract): Geological Society of America Special Paper, v. 115, p. 351.

———1970, Fumarolic mounds and ridges of the Bishop Tuff, California: Geological Society of America Bulletin, v. 81, p. 851–868.

Sieh, Kerry, and Bursik, Marcus, 1986, Most recent eruption of the Mono Craters, eastern central California: Journal of Geophysical Research v. 91, no. B12, p. 12,539–12,571.

Smith, J.L., and Rex, R.W., 1977, Drilling results from eastern Long Valley caldera: American Nuclear Society Meeting on Energy and Mineral Recovery Research, Golden, Colorado, April 12–14, 1977, p. 529–540.

Smith, R.L., and Bailey, R.A., 1968, Resurgent cauldrons: Geological Society of America Memoir 116, p. 613–662.

Sorey, M.L., 1978, The hydrothermal system of Long Valley caldera, California: U.S. Geological Survey Professional Paper 1044-A, p. A1–A60.

———1985, Evolution and the present state of the hydrothermal system in Long Valley caldera: Journal of Geophysical Research, v. 90, no. B13, p. 11,219–11,228.

Steeples, D.W., and Iyer, H.M., 1976, Low-velocity zone under Long Valley as determined from teleseismic events: Journal of Geophysical Research, v. 81, no. 5, p. 849–860.

Stine, Scott, 1984, Late Holocene lake level fluctuations and island volcanism at Mono Lake, California, *in* Holocene paleoclimatology and tephrochronology east and west of the central Sierran Crest, Field Trip Guidebook for the Friends of the Pleistocene, Pacific Cell, October 12–14, 1984, p. 21–49.

Suemnicht, Gene, 1987, Results of deep drilling in the western moat of Long Valley, California: EOS, American Geophyiscal Union Transactions, v. 68, no. 40, p. 785, 798.

Van Wormer, J.D., and Ryall, A.S., 1980, Sierra Nevada-Great Basin boundary zone: earthquake hazard related to structure, active tectonic processes, and anomalous patterns of earthquake occurrence: Seismological Society of America Bulletin, v. 70, p. 1557–1572.

Wollenberg, H.A., Sorey, M.L., Ferrar, C.D., White, A.F., Flexer, Steve, and Bartel, L.C., 1987, A core hole in the southwestern moat of Long Valley caldera: early results: EOS, American Geophyiscal Union Transactions, v. 68, no. 20, p. 529–534.

Wood, S.H., 1977, Distribution, correlation, and radiocarbon dating of late Holocene tephra, Mono and Inyo Craters, eastern California: Geological Society of America Bulletin, v. 88, p. 89–95.

———1983, Chronology of late Pleistocene and Holocene volcanics, Long Valley and Mono Basin geothermal areas, eastern California: U.S. Geological Survey Open-File Report 83–747, 76 p.

☆U.S. Government Printing Office : 1989 - 242-384/05810

best copy available